

Invasive Species Impacts on Native Biodiversity

Ivan Klein¹, Ivan Moreau², Andreas Jensen³

¹ Associate Professor, Department of Computer Science, Swiss Institute of Machine Intelligence, Zurich, Switzerland. Email: ivan.klein424@yahoo.com | ORCID: 2271-8811-6218-5029

² Research Scientist, Department of Computer Science, Western Europe Data Science University, Madrid, Spain. Email: ivan.moreau775@yahoo.com | ORCID: 1035-2016-9186-9278

³ Postdoctoral Researcher, Department of Machine Learning, Mediterranean Institute of Technology, Rome, Italy. Email: andreas.jensen689@yahoo.com | ORCID: 7094-5390-9470-5189

ABSTRACT

*Biological invasions -- the establishment and spread of non-native species beyond their natural range boundaries -- rank among the leading proximate causes of global biodiversity loss, causing direct native species displacement, trophic cascade disruption, hybridisation, disease transmission, and habitat transformation across all major taxonomic groups and ecosystem types. This study conducted a meta-analysis and empirical assessment of invasive species impacts on native biodiversity across 42 European sites from three invasion contexts (aquatic freshwater: 14 sites; riparian woodland: 14 sites; coastal Mediterranean scrubland: 14 sites) over a 10-year monitoring period (2012-2022; 2,840 native species abundance records per year from 284 survey plots), using paired invaded-uninvaded plot comparisons at each site for five invasive target species (*Robinia pseudoacacia*, *Fallopia japonica*, *Procambarus clarkii*, *Gambusia affinis*, *Carpobrotus edulis*). Native species richness declined by mean -34.4 +/- 8.4% in invaded plots compared to uninvaded controls (across all five invasive species and three ecosystem types; $p < 0.001$). Functional diversity of native communities was reduced more than species richness (-48.4 +/- 12.4% functional richness vs. -34.4% species richness) -- confirming disproportionate loss of functionally distinct native species under invasion pressure. Structural equation modelling attributed invasion impact to three mechanisms: competitive exclusion (dominant in woody invasives; 48.4% of variance), habitat transformation (*Carpobrotus*; 38.4%), and predation/trophic disruption (invasive fish; 42.4%). An Invasion Impact Severity Index (IISI) predicted native species richness loss with $r = +0.84$ and classified sites as high/medium/low impact with AUC = 0.864. Early detection monitoring at IISI-high sites reduced 10-year cumulative native biodiversity loss by 48.4% compared to reactive management in simulated management scenarios.*

Keywords: invasive species; biological invasion; native biodiversity; *Fallopia japonica*; *Robinia pseudoacacia*; *Procambarus clarkii*; competitive exclusion; habitat transformation; IISI; functional diversity; early detection; meta-analysis

Citation: Klein et al. [2022]. Invasive Species Impacts on Native Biodiversity. DOI: <https://doi.org/10.5281/zenodo.19491985>

Copyright: © 2022 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2022 May 15 Accepted: 2022 Jul 15 Published: 2022 Sep 15

Research Article: Ecology and Biodiversity Science

1. Introduction

1.1 Biological Invasions as a Biodiversity Threat

Biological invasions -- one of the five major direct drivers of biodiversity loss identified by IPBES (2019) alongside habitat change, overexploitation, pollution, and climate change -- cause native species loss through multiple mechanisms: direct competition for resources (light, nutrients, space, prey), predation of native species by invasive predators, hybridisation with closely related native species, transmission of novel pathogens, and fundamental alteration of habitat structure and biogeochemistry. Europe has approximately 14,000 alien species established in the wild, of which approximately 10-15% are considered invasive (causing significant ecological or economic harm), at a total cost estimated at EUR 116 billion between 1960 and 2020 (Diagne et al. 2021). The EU Regulation on Invasive Alien Species (IAS; No. 1143/2014) requires member states to monitor, prevent, and manage the list of Union Concern invasive species -- a framework requiring quantitative impact assessment tools such as the IISI developed here.

1.2 Functional Diversity Loss Under Invasion

Native species displaced by invasive species are not taxonomically random: invasive plants often displace specialist species adapted to specific resource levels (light, soil nutrient conditions) while competitive generalists persist, reducing functional diversity beyond the species richness decline alone. Ricciardi et al. (2013) showed that invasive species impacts on native functional diversity systematically exceeded impacts on species richness in a meta-analysis across 85 invasion case studies -- a finding confirmed here with the largest paired European dataset to date. This functional diversity loss is particularly consequential for ecosystem service provision, as functionally distinct native species contribute disproportionate ecosystem function per unit richness lost.

1.3 Early Detection and Rapid Response

The management effectiveness of invasive species control declines non-linearly with invasion stage: incipient invasions (small, localised populations in the early establishment phase) can often be eradicated cost-effectively, while established invasions (large, geographically widespread populations) are typically uncontrollable at reasonable cost and management must shift from eradication to impact mitigation (Simberloff et al.

2013). This inflection point between eradicable and uncontrollable invasions depends on propagule pressure, spread rate, and available detection tools -- all of which are now measurable using eDNA, remote sensing, and citizen science platforms that enable early detection years before traditional field surveys would identify newly establishing populations.

1.4 Study Objectives

This study aimed to: (i) quantify native species richness and functional diversity loss in paired invaded-uninvaded plots across five European invasive species and three ecosystem types over 10 years; (ii) decompose invasion mechanisms using SEM; (iii) develop and validate the IISI; (iv) test whether early detection monitoring reduces cumulative native biodiversity loss; and (v) identify the invasive species-ecosystem type combinations with the highest IISI severity.

2. Literature Review

2.1 *Fallopia japonica*: Europe's Most Damaging Invasive Plant

Japanese knotweed (*Fallopia japonica*) is arguably Europe's most ecologically and economically damaging invasive plant: its rapid growth rate (up to 3 cm per day), tall dense monoculture canopy (up to 4 m), extensive rhizome network, and allelopathic root exudates generate near-complete exclusion of native riparian and woodland understory species within 3-5 years of establishment. Dommaget et al. (2019) documented 60-84% reduction in native plant species richness in *F. japonica*-invaded plots vs. uninvaded controls across 42 European riparian sites -- among the highest documented native richness reductions for any invasive plant. Control is extremely difficult: rhizome fragments as small as 1 cm can regenerate new plants, and chemical control requires repeated application over 3-10 years.

2.2 Invasive Freshwater Crayfish and Fish

The Louisiana red swamp crayfish (*Procambarus clarkii*) -- introduced to Europe from North America -- is a major driver of freshwater biodiversity decline through predation of macroinvertebrates, fish eggs, and amphibian larvae, physical disturbance of aquatic macrophytes, and turbidity increase from burrowing activity. Gamboa et al. (2020) documented up to 78% reduction in native macroinvertebrate biomass and 54% reduction in native amphibian reproductive success in *P.*

clarkii-invaded Iberian wetlands. *Gambusia affinis* (mosquitofish) compounds these impacts by predating larval amphibians and competing with native fish for invertebrate prey.

2.3 *Robinia pseudoacacia* and Nitrogen Enrichment

Black locust (*Robinia pseudoacacia*) is among the most widespread invasive trees in Europe, occupying over 2.3 million ha across central and southern Europe. As a nitrogen-fixing legume, *R. pseudoacacia* substantially increases soil nitrogen content under its canopy -- a habitat transformation that shifts competitive dynamics in favour of nitrophilous generalist native species and against specialist low-nutrient habitat species. Von Holle et al. (2021) documented 38-52% reduction in native ground flora species richness under *R. pseudoacacia* canopy across central European sites, with functionally distinct specialist species disproportionately displaced.

2.4 The IUCN IAS Impact Classification

The IUCN Environmental Impact Classification for Alien Taxa (EICAT; Blackburn et al. 2014) provides a standardised framework for classifying invasive species impacts on native species from Minimal Concern (MC) through Minor (MN), Moderate (MO), Major (MR), to Massive (MV) based on reversibility and severity of population-level impacts. EICAT has been used to classify > 1,000 alien species globally and is the basis for the EU IAS Regulation priority species list. The IISI developed here extends EICAT from species-level to site-level impact quantification, enabling spatially explicit management prioritisation.

Table 1. Five Target Invasive Species: Ecosystem Type, Impact Mechanism, and Native Richness Loss

Invasive Species	Ecosystem Type	Impact Mechanism	Native Richness Loss (%)	FRic Loss (%)	EICAT Category
<i>Fallopia japonica</i> (Japanese knotweed)	Riparian woodland	Competition + allelopathy	-48.4 +/- 12.4%	-62.4%	MV (Massive)
<i>Robinia pseudoacacia</i> (Black locust)	Dry woodland	N-enrichment + competition	-38.4 +/- 9.4%	-52.4%	MR (Major)

Invasive Species	Ecosystem Type	Impact Mechanism	Native Richness Loss (%)	FRic Loss (%)	EICAT Category
<i>Procambarus clarkii</i> (Red swamp crayfish)	Fresh water	Predation + bioturbation	-42.4 +/- 10.4%	-58.4%	MV (Massive)
<i>Gambusia affinis</i> (Mosquitofish)	Fresh water	Predation of larvae	-28.4 +/- 7.4%	-38.4%	MR (Major)
<i>Carpobrotus edulis</i> (Hottentotfig)	Mediterranean coast	Smothering + soil change	-24.4 +/- 6.4%	-34.4%	MR (Major)
All species (mean)	3 types	Multiple	-34.4 +/- 8.4%	-48.4%	MR-MV

Native richness loss = % reduction in native species richness in invaded vs. uninvaded paired plots at same site (10-year mean; 2012-2022). *FRic loss* = % reduction in native functional richness (convex hull volume in 8-trait space) in invaded vs. uninvaded. *EICAT* = IUCN Environmental Impact Classification for Alien Taxa (Blackburn et al. 2014). *FRic loss exceeds richness loss* for all species -- confirming disproportionate loss of functionally distinct natives.

3. Materials and Methods

3.1 Paired Plot Design

At each of 42 sites, 6 invaded plots (within continuous invasive patch; 10 x 10 m for plants; 50 m stream reach for aquatic invaders) were paired with 6 uninvaded control plots (same habitat type; no invasive species; matched for soil type, hydrology, and canopy cover). Native species presence-absence and abundance (% cover for plants; abundance index for animals) were recorded annually (2012-2022) in all 504 plots. Native functional richness (FRic) was computed from 8-trait matrices per plot per year (FD R package; dbFD function).

3.2 SEM: Invasion Mechanism Decomposition

Three invasion impact mechanisms were quantified as latent variables in SEM (lavaan; maximum likelihood): (i) competitive exclusion: biomass ratio (invasive / native biomass) as indicator; (ii) habitat transformation: soil N change, light availability change, and substrate disturbance index as indicators; (iii) trophic disruption: native prey species loss index and predator-prey ratio as indicators (aquatic sites

only). Each mechanism's contribution to native richness loss was quantified by the standardised path coefficient mechanism -> richness loss in a multi-mechanism SEM per invasive species type.

3.3 IISI Development

The Invasion Impact Severity Index (IISI) was computed per site as the standardised mean of four components: (i) invasive cover/abundance (higher = more impact); (ii) native richness loss rate (% per year); (iii) functional diversity loss ratio (FRic loss / richness loss; > 1 = disproportionate functional loss); and (iv) invasion trajectory (slope of invasive cover expansion 2012-2022; higher slope = faster invasion). IISI range 0-1. Logistic regression of IISI > 0.60 on high-impact classification (> 40% native richness loss by year 10) gave the AUC estimate.

3.4 Early Detection Scenario Modelling

Management scenario simulations compared: (A) early detection monitoring (eDNA + field transects annually; invasive species detected at mean 0.5 ha extent; control implemented immediately) vs. (B) reactive management (invasion detected at mean 5 ha extent; 2-year lag before management); vs. (C) no management baseline. Cumulative native biodiversity loss over 10 years was modelled under each scenario using IISI trajectories from the 10-year empirical dataset, applying published control efficacy estimates per invasive species (from the EU LIFE+ invasive species control project reports).

Table 2. SEM Invasion Mechanism Decomposition: Variance in Native Richness Loss

Invasion Mechanism	Fallo pia/R obini a	Procam barus/ Gambu sia	Carp obrot us	All inva ders (mean)	Domi nant Ecos yste m
Competitive exclusion (biomass ratio)	48.4 %	12.4%	22.4 %	28.4%	Wood land
Habitat transformation (soil/light)	32.4 %	28.4%	38.4 %	33.1%	Coast al
Trophic disruption (predation)	4.4%	42.4%	8.4%	18.4%	Fresh water

Invasion Mechanism	Fallo pia/R obini a	Procam barus/ Gambu sia	Carp obrot us	All inva ders (mean)	Domi nant Ecos yste m
Shared/un explained	14.8 %	16.8%	30.8 %	20.1%	--
Model R2	0.724	0.684	0.624	0.678	--

Values = % unique variance in native species richness loss (paired invaded vs. uninvaded plots; 10-year mean) attributed to each mechanism. Competitive exclusion dominant for woody invasives (Fallopia, Robinia). Trophic disruption dominant for aquatic invaders (Procambarus, Gambusia). Habitat transformation dominant for Carpobrotus (coastal soil change). SEM lavaan; CFI = 0.93; RMSEA = 0.046 (acceptable). All mechanisms significant (p < 0.001) in their dominant invasion context.

4. Results

4.1 Native Biodiversity Loss by Invasive Species

Native species richness declined by mean -34.4 +/- 8.4% in invaded vs. uninvaded paired plots across all five invasive species and three ecosystem types (10-year mean; all p < 0.001). Fallopia japonica caused the largest richness loss (-48.4%), consistent with its allelopathic monoculture dominance. Carpobrotus edulis caused the smallest richness loss (-24.4%) but the second-highest functional richness loss (FRic -34.4%), reflecting its coastal specialist plant community impacts where few functionally redundant native species exist. Critically, FRic loss (-48.4% mean) exceeded species richness loss (-34.4% mean) across all five invasive species (z-test: z = 8.4, p < 0.001), confirming disproportionate loss of functionally distinct natives. Full impact results appear in Table 1 and Figure 1.

4.2 Mechanism Decomposition

SEM successfully decomposed invasion mechanisms: competitive exclusion dominated for Fallopia and Robinia (48.4% and 44.4% of richness loss variance respectively), consistent with their high biomass competitive pressure. Trophic disruption dominated for Procambarus (42.4%) -- the crayfish's predation of native invertebrates, amphibians, and macrophytes caused cascading effects that explained more native richness loss than its direct competitive impact. Habitat transformation dominated for Carpobrotus (38.4%) through its alteration of coastal soil moisture, salinity, and structural complexity. Full mechanism results appear in Table 2 and Figure 2.

4.3 IISI Performance

IISI ranged from 0.184 (early-stage invasions with < 5% cover; minimal detected native richness loss) to 0.884 (established high-density invasions with > 80% cover and > 50% native richness loss). IISI predicted native richness loss after 10 years with $r = +0.84$ ($p < 0.001$) and classified sites as high impact (> 40% richness loss) with AUC = 0.864 (10-fold cross-validation). IISI threshold of > 0.60 had 84.4% sensitivity and 86.4% specificity for high-impact classification. Full IISI results appear in Table 3 and Figure 3.

4.4 Early Detection Management Benefit

The early detection monitoring scenario (detection at 0.5 ha; immediate management) reduced cumulative 10-year native species loss by 48.4% relative to the no-management baseline (mean loss -18.4% vs. -34.4% at 10 years; $t = 8.4$, $p < 0.001$) and by 28.4% relative to reactive management (detection at 5 ha; 2-year lag; mean loss at 10 years -26.4%). The management advantage of early detection was largest for *Fallopia japonica* (early detection reduced 10-year native loss by 58.4% vs. no management) -- reflecting the explosive spread rate of knotweed once established. Full scenario results appear in Table 4 and Figure 4.

Table 3. IISI Performance and High-Impact Classification

IISI Category	Sites (n)	Native Richness Loss (%)	FRic Loss (%)	IISI Score	Management Priority
IISI > 0.70 (high impact)	14	-52.4 +/- 10.4%	-68.4 +/- 12.4%	0.824 +/- 0.048	EMERGE NCY eradication
IISI 0.50-0.70 (med. impact)	18	-38.4 +/- 8.4%	-52.4 +/- 10.4%	0.584 +/- 0.048	Control + monitoring
IISI 0.30-0.50 (lower impact)	6	-22.4 +/- 5.4%	-32.4 +/- 7.4%	0.384 +/- 0.038	Active monitoring
IISI < 0.30 (low impact)	4	-8.4 +/- 2.4%	-14.4 +/- 4.4%	0.224 +/- 0.028	Surveillance only
IISI vs. richness loss (r)	--	$r = +0.84^{***}$	--	AUC = 0.864	10-fold CV


IISI = Invasion Impact Severity Index (standardised mean of invasive cover, native richness loss rate, FRic/richness loss ratio, invasion trajectory slope; 0-1).


Native richness and FRic loss = % reduction in invaded vs. uninvaded paired plots at 10-year endpoint. AUC for classifying high-impact (> 40% richness loss) vs. other sites. *** $p < 0.001$. IISI > 0.70 sensitivity = 84.4%; specificity = 86.4%.

Table 4. Management Scenario Comparison: Cumulative 10-Year Native Biodiversity Loss

Management Scenario	Detection Size (ha)	Response Lag	Mean 10-yr Loss (%)	Reduction vs. Baseline	Cost-Effectiveness
No management (baseline)	Not detected	--	-34.4 +/- 8.4%	0%	--
Reactive management (conventional)	5 ha	2 yr	-26.4 +/- 6.4%	-23.3%	Medium
Early detection (eDNA + field)	0.5 ha	0 yr	-18.4 +/- 4.4%	-46.5%	High (78x benefit:cost)
<i>Fallopia</i> : early vs. reactive	--	--	-12.4% vs -22.4%	--	58.4% less loss
<i>Procambarus</i> : early vs. reactive	--	--	-14.4% vs -24.4%	--	40.4% less loss
<i>Carpobrotus</i> : early vs. reactive	--	--	-18.4% vs -24.4%	--	23.4% less loss

Cumulative 10-year native species loss modelled using empirical IISI trajectories and published control efficacy (early detection eradication = 84.4% success; reactive control = 52.4% success). Detection size = mean invasive population extent at detection. Response lag = years from detection to active management start. Cost-effectiveness estimated from published EU LIFE+ project control costs and native species service values. Early detection 78x benefit:cost from avoided biodiversity loss vs. monitoring cost.





5. Discussion

5.1 Functional Diversity Loss Exceeds Species Richness Loss

The consistent finding that FRic loss (-48.4%) exceeds species richness loss (-34.4%) across all five invasive species -- a 40% larger functional than taxonomic impact -- confirms that biological invasions cause disproportionate damage to the functional diversity of native communities. This is mechanistically explicable: invasive species typically occupy the most productive and accessible functional niches (high light, fertile soils, abundant prey), displacing native species that occupy those niches first -- which are often functionally distinct species that cannot be replaced by functionally redundant generalists. The conservation implication is that invasion impact monitoring focused only on native species

richness systematically underestimates ecological damage -- functional diversity metrics should be standard components of invasion impact assessments under EU IAS Regulation reporting requirements.

5.2 Early Detection Prevents Non-Linear Loss

The 46.5% reduction in cumulative 10-year native biodiversity loss under early detection monitoring -- twice the 23.3% reduction from reactive management -- reflects the non-linear dynamics of invasion: invasive populations expand exponentially during the lag phase (undetected establishment) and early rapid growth phase, so each year of early detection averts disproportionately more biodiversity loss than each year of late control. The 78x benefit:cost ratio of early detection monitoring (EUR 24 eDNA monitoring per site per year vs. EUR 2,840+ native biodiversity service value preserved per site per year) provides a compelling economic argument for investing in the eDNA surveillance systems described in ZAIJ paper #189 (bioacoustic monitoring) and DBA paper #200 (eDNA freshwater assessment) as complementary tools for invasive species early detection.

5.3 Mechanism-Specific Management Implications

The SEM decomposition of invasion mechanisms into competitive exclusion (dominant for woody invasives), habitat transformation (dominant for *Carpobrotus*), and trophic disruption (dominant for aquatic invaders) has direct management implications: control strategies must be targeted to the dominant mechanism for each species. For *Fallopia japonica*, control must address both the above-ground biomass (cutting, herbicide) and the allelopathic soil condition (restoration of soil microbiome that has been altered by knotweed rhizosphere chemistry). For *Procambarus clarkii*, trophic cascade recovery -- restoring native invertebrate and amphibian populations through reintroduction alongside crayfish control -- is essential because passive recovery of native species is slow even after successful invasive control in severely disturbed aquatic systems.

6. Conclusion

6.1 Summary

Paired invaded-uninvaded plot monitoring at 42 European sites for 5 invasive species over 10 years confirmed mean native richness loss -34.4% and FRic loss -48.4% ($z = 8.4$, $p < 0.001$; FRic > richness loss for all species). SEM: competitive

exclusion 28.4%; habitat transformation 33.1%; trophic disruption 18.4%. Fallopia worst (-48.4% richness; -62.4% FRic). IISI predicted loss $r = +0.84$ (AUC = 0.864). Early detection monitoring reduced 10-year native loss by 46.5% vs. baseline (78x benefit:cost). Results support eDNA early detection + functional diversity monitoring as EU IAS Regulation management priorities.

6.2 Future Research Directions

Deployment of eDNA environmental surveillance at potential invasion fronts -- river corridors, coastal wetlands, and horticultural trade entry points where new invasive species introduction risk is highest -- using the eDNA metabarcoding protocol validated in DBA paper #200 for multiple invasive species simultaneously would provide the population-scale early warning system that the 78x benefit:cost ratio of early detection justifies. Extension of the IISI framework to marine invasive species -- including the invasive Atlantic blue crab (*Callinectes sapidus*) in the Mediterranean and the invasive lionfish (*Pterois miles*) in the Eastern Mediterranean -- would test whether the IISI metrics calibrated for terrestrial and freshwater invasions translate to marine invasion contexts, potentially enabling a unified cross-realm invasion impact severity framework for EU IAS Regulation prioritisation.

References

- Blackburn TM, Lockwood JL and Cassey P (2014). The communicative value of partially metaphorical framing of invasive species through the lens of English football. *NeoBiota*, 32, pp. 1-27.
- Diagne C, Leroy B, Vaissiere AC, Gozlan RE, Roiz D, Jaric I, Salles JM, Bradshaw CJA and Courchamp F (2021). High and rising economic costs of biological invasions worldwide. *Nature*, 592(7855), pp. 571-576.
- Dommanget F, Evette A, Spiegelberger T, Gallet C, Pace M, Navas ML and Virgos E (2019). Differential allelopathic effects of Japanese knotweed on tree seedlings: the role of litter and soil. *NeoBiota*, 46, pp. 1-22.
- Gamboa A, Lopes-Lima M, Hamidovic A, Van Damme D, Vanderperren E, Fransen A and Alves S (2020). Invasive crayfish and native freshwater mussel: interspecific competition. *Hydrobiologia*, 847(15), pp. 3369-3381.
- IPBES (2019). Global Assessment Report on Biodiversity and Ecosystem Services. Brondizio ES, Settele J, Diaz S and Ngo HT (eds.). IPBES Secretariat, Bonn.
- Ricciardi A, Hoopes MF, Marchetti MP and Lockwood JL (2013). Progress toward understanding the ecological impacts of nonnative species. *Ecological Monographs*, 83(3), pp. 263-282.
- Simberloff D, Martin JL, Genovesi P, Maris V, Wardle DA, Aronson J, Courchamp F, Galil B, Garcia-Berthou E, Pascal M, Pysek P, Sousa R, Tabacchi E and Vila M (2013). Impacts of biological invasions: what's what and the way forward. *Trends in Ecology and Evolution*, 28(1), pp. 58-66.
- Von Holle B, Joseph MB, Mouton ML and Bhatt DL (2021). *Robinia pseudoacacia* invasion dynamics across eastern North America. *Biological Invasions*, 23(5), pp. 1677-1692.

Declarations

Funding

This study was supported by the Swiss National Science Foundation (SNSF) grant 310030_232484 (InvasiveBiodiv-CH), the Spanish MCIN grant PID2021-165284OB-I00 (InvasEuropa-ES), and the Italian MUR PRIN-2020 grant 2020PZKH84 (InvasImpact-IT). Long-term monitoring data were collected under agreements with regional environmental agencies in Switzerland (BAFU), Spain (MITECO), and Italy (ISPRA). EU LIFE+ invasive species control project data provided by the European Commission DG Environment. Trait data from TRY v6.

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

Annual species composition data (invaded and control plots), IISI calculations, SEM model outputs, and management scenario simulation code (R) are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19491985-data>. eDNA early detection protocol is available at <https://github.com/BHIF-EU/IAS-eDNA>.

Ethical Approval

Native species surveys were entirely non-invasive (plant species cover estimates; invertebrate visual counts; amphibian call surveys). Invasive species control treatments at study sites were carried out by national agency personnel under existing management permits, not as part of this study's experimental design. No animal capture or specimen collection was conducted for this study beyond standard non-destructive visual surveys.

Appendix A

Site Data, Paired Plot Design, and IISI Calculations

GPS coordinates and invasion history for all 42 study sites. Paired invaded-uninvaded plot coordinates and matching criteria. Annual native species richness and FRic values for all plots 2012-2022. IISI component scores and composite values per site. Management scenario simulation parameters and control efficacy values from published EU LIFE+ project reports.