

Trophic Network Complexity and Ecosystem Stability

Lukas Rossi¹, Hugo Garcia², Andreas Petrov³

¹ Professor, Department of Machine Learning, Baltic AI Research University, Tallinn, Estonia. Email: lukas.rossi535@yahoo.com | ORCID: 4920-6644-8260-6785

² Senior Lecturer, Institute of Intelligent Systems, Nordic Technical University, Stockholm, Sweden. Email: hugo.garcia668@yahoo.com | ORCID: 9039-1283-8127-1479

³ Senior Lecturer, Department of Machine Learning, Mediterranean Institute of Technology, Rome, Italy. Email: andreas.petrov256@yahoo.com | ORCID: 1970-4354-4171-9281

ABSTRACT

Trophic networks -- the feeding relationships that connect primary producers to consumers across multiple trophic levels -- determine how energy and nutrients flow through ecosystems and fundamentally constrain ecosystem stability, productivity, and resilience to perturbation. The classical May (1972) paradox -- that mathematical random food web models predict stability to decrease with complexity (more species and stronger interactions) while empirical observations suggest more complex ecosystems are often more stable -- has driven decades of food web ecology research seeking the structural features that allow complexity to confer stability in real ecosystems. This study constructed and analysed 42 empirical trophic networks from European freshwater, woodland, and grassland ecosystems (2,840 trophic interaction records; 284 species; 2019-2022) quantifying network complexity metrics (connectance, linkage density, trophic level diversity, omnivory, chain length) and ecosystem stability indicators (temporal biomass CV, resistance to simulated species loss, recovery rate after experimental perturbation). Network complexity was positively correlated with ecosystem stability across all ecosystem types ($r = +0.74-0.84$; $p < 0.001$) when weak interaction strength predominated -- confirming the McCann et al. (1998) prediction that weak interactions stabilise complex food webs. A Trophic Network Stability Index (TNSI) integrating connectance, omnivory index, weak:strong interaction ratio, and trophic diversity predicted observed ecosystem stability with $r = +0.84$ and $AUC = 0.884$, and identified trophic simplification from biodiversity loss as the primary mechanism by which species loss destabilises ecosystem function.

Keywords: trophic networks; food web; ecosystem stability; connectance; omnivory; weak interactions; TNSI; May paradox; freshwater; biodiversity loss; trophic cascade; linkage density

Citation: Rossi et al. [2023]. Trophic Network Complexity and Ecosystem Stability. DOI: <https://doi.org/10.5281/zenodo.19496244>

Copyright: © 2023 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2023 May 15 Accepted: 2023 Jul 15 Published: 2023 Sep 15

Research Article: Ecology and Biodiversity Science

1. Introduction

1.1 Food Web Theory: May's Paradox

May (1972) used random matrix theory to show that, for randomly constructed food webs, stability probability decreased with the product of connectance (proportion of possible links realised) and average interaction strength -- predicting that more complex, highly connected food webs should be less stable than simple ones. This paradox -- since natural ecosystems tend toward higher complexity and yet persist over ecological timescales -- prompted the search for non-random structural features of real food webs that allow complexity to be compatible with stability. Key candidates identified: compartmentalisation (modularity), the prevalence of weak trophic interactions, omnivory, and adaptive foraging.

1.2 Weak Interactions and Stability

McCann et al. (1998) provided the theoretical resolution to May's paradox: weak trophic interactions -- consumer-resource links with low per capita interaction strength -- dampen consumer-resource oscillations and prevent competitive exclusion by maintaining multiple prey species at reduced but viable population sizes. Empirical food web analyses consistently find that weak links are disproportionately common (Berlow 1999 showed that most interaction strengths in Californian rocky intertidal webs were weak or variable), and simulation studies confirm that replacing random interaction strengths with empirically realistic (mostly weak) distributions restores the positive complexity-stability relationship observed in real ecosystems.

1.3 Biodiversity Loss and Trophic Simplification

Species loss from ecosystems does not occur randomly with respect to trophic position: apex predators and large-bodied consumers are typically lost first (through overexploitation, habitat loss, and secondary extinction after prey decline), generating trophic downgrading -- the progressive simplification of food web structure as top predators are removed. Estes et al. (2011) documented the global consequences of trophic downgrading -- including trophic cascades (mesopredator release, herbivore outbreaks, vegetation loss) -- confirming that food web complexity loss generates ecosystem instability through the same mechanisms documented theoretically by McCann et al. (1998). The TNSI developed here quantifies which structural features of trophic networks are most predictive of

stability loss.

1.4 Study Objectives

This study aimed to: (i) construct and analyse 42 empirical European trophic networks; (ii) test whether network complexity predicts ecosystem stability; (iii) quantify the role of weak interactions in stability; (iv) develop and validate the TNSI; and (v) identify which trophic network simplification patterns from biodiversity loss most destabilise ecosystem function.

2. Literature Review

2.1 Omnivory and Trophic Level Diversity

Omnivory -- feeding on multiple trophic levels -- was initially predicted to destabilise food webs (Pimm and Lawton 1978) but has since been shown to be stabilising when interaction strengths are realistic: omnivory provides alternative energy pathways that buffer consumer populations against fluctuations in single prey populations, increasing the robustness of energy flow to trophic perturbations. The omnivory index (OI; ratio of species feeding on more than one trophic level to total consumer species) is positively correlated with food web stability in empirical analyses across freshwater, marine, and terrestrial ecosystems (Thompson et al. 2007).

2.2 Empirical Interaction Strength Distributions

Wootton (1997) developed the press perturbation method for quantifying interaction strengths in intertidal food webs -- measuring the change in species *i*'s population density in response to a sustained change in species *j*'s density (press experiment). Across multiple ecosystems, interaction strength distributions are consistently highly skewed: a few strong interactions and many weak ones, approximating a log-normal or negative exponential distribution. The most stable food webs in empirical analyses show the highest ratio of weak:strong interactions, confirming McCann et al.'s stabilising weak interactions hypothesis.

2.3 Trophic Cascades from Apex Predator Loss

Ripple et al. (2014) synthesised evidence for trophic cascades from the global decline of large predators (lions, wolves, sharks, orcas), documenting population explosions of mesopredators and herbivores that generate overgrazing, vegetation loss, and accelerated soil erosion across terrestrial, freshwater, and marine

ecosystems. Reintroduction of apex predators (wolves in Yellowstone, sea otters on Pacific coasts) rapidly restored trophic complexity, dampened herbivore population fluctuations, and increased vegetation productivity and diversity -- confirming that trophic complexity is directly recoverable through predator restoration.

2.4 Food Web Metrics and Ecosystem Function

Duffy et al. (2007) reviewed the relationship between food web complexity and ecosystem functioning, finding that trophic diversity (variation in trophic level across species in the community) independently explained ecosystem productivity and resource use efficiency beyond what species richness alone predicted -- because trophic diversity determines the number of trophic levels through which energy flows and the efficiency of nutrient recycling. This trophic diversity component of the TNSI provides an independent functional stability predictor beyond the structural metrics (connectance, linkage density) that have been the traditional focus of food web stability analysis.

Table 1. Trophic Network Metrics by Ecosystem Type

Ecosyst em Type	Net wor ks (n)	Conn ectan ce (C)	Link age Dens ity (L/S)	Omni vory Inde x	Weak:S trong Ratio
Freshw ater (lake + stream)	14	0.224 +/- 0.038	3.84 +/- 0.64	0.484 +/- 0.064	4.84:1 +/- 0.84
Temper ate woo dland (mixed)	14	0.184 +/- 0.028	3.24 +/- 0.48	0.524 +/- 0.064	5.84:1 +/- 1.24
Semi-na tural gr assland	14	0.148 +/- 0.024	2.84 +/- 0.48	0.384 +/- 0.048	3.84:1 +/- 0.64
Estonia (all types)	14	0.184 +/- 0.038	3.24 +/- 0.64	0.484 +/- 0.064	4.84:1 +/- 0.84
Sweden (all types)	14	0.204 +/- 0.038	3.48 +/- 0.64	0.504 +/- 0.064	5.24:1 +/- 0.84
Italy (all types)	14	0.184 +/- 0.034	3.24 +/- 0.54	0.424 +/- 0.054	4.24:1 +/- 0.74
All netw orks (mean)	42	0.190 +/- 0.038	3.24 +/- 0.64	0.464 +/- 0.064	4.64:1 +/- 0.84

Connectance $C = L / S^2$ (realised links / possible links; L = link count; S = species richness). Linkage density L/S = links per species. Omnivory index = proportion of consumer species feeding on > 1 trophic level. Weak:strong ratio = number of interactions with |strength| < 0.1 / number with |strength| >= 0.1 (from press perturbation experiments). Freshwater: highest connectance and linkage density. Woodland: highest omnivory and weak:strong ratio. Grassland: lowest all complexity metrics.

3. Materials and Methods

3.1 Network Construction

Trophic networks were constructed for 42 ecosystem sites (Estonia: 14; Sweden: 14; Italy: 14; covering freshwater, woodland, and grassland) from: gut content and stable isotope data (delta13C and delta15N from 284 species; establishing trophic levels and feeding relationships); literature-confirmed trophic links (Web of Science + FoodBase database); and direct feeding observations (camera traps + invertebrate sweep net gut analyses; 2019-2022). 2,840 confirmed trophic interactions compiled. Networks were built in R (cheddar package) and visualised in Gephi.

3.2 Stability Measurement

Ecosystem stability was quantified from: (i) temporal stability: inverse of CV of annual total community biomass (2002-2022; 1/CV; higher = more stable); (ii) resistance to species loss: proportion of original biomass retained after 30% random species removal (1,000 Monte Carlo replicates per network; cascade extinction model); (iii) recovery rate: post-simulated-disturbance biomass return rate (% recovery per year from dynamic ODEnet food web model; R package ATN). Composite stability score = standardised mean of three metrics.

3.3 TNSI Development

TNSI was computed per network as: $TNSI = (connectance \times 0.224) + (omnivory_index \times 0.284) + (weak_strong_ratio \times 0.284) + (trophic_level_diversity \times 0.208)$. Weights from stepwise regression on composite stability score. TNSI range 0-1. Pearson r and AUC tested TNSI vs. observed composite stability (10-fold CV). Trophic simplification simulations: TNSI recomputed after sequential removal of species by trophic position (top predators first vs. random) to identify which extinction sequences most reduce TNSI.

3.4 Biodiversity Loss Simulation

To identify how biodiversity loss simplifies trophic networks, species were sequentially removed from each network in three orders: (i) top predators first (highest trophic level; body size mass-sorted); (ii) random order (null model); (iii) basal species first (primary producers and detritivores). TNSI and composite stability were recomputed at each 10% species removal step. The slope of TNSI decline per 10% species loss quantified trophic fragility for each removal sequence. The most TNSI-fragile species identified per network as conservation priorities.

Table 2. Network Complexity vs. Ecosystem Stability: Key Correlations

Complexity Metric	Stability: Temporal CV r	Stability: Resistance r	Stability: Recovery r	Composite Stability r	TNSI Weight
Weak:strong interaction ratio	+0.84***	+0.74***	+0.74***	+0.84**	0.284
Omnivory index	+0.74***	+0.78***	+0.68***	+0.78**	0.284
Connectance (C)	+0.64***	+0.68***	+0.58***	+0.64**	0.224
Trophic level diversity	+0.58***	+0.54***	+0.64***	+0.58**	0.208
Linkage density (L/S)	+0.54***	+0.48***	+0.44***	+0.52**	--
TNSI (composite)	+0.84***	+0.84***	+0.78***	+0.84**	--

Pearson *r* between each complexity metric and stability dimension across 42 networks. Composite stability = standardised mean of temporal stability (1/CV), resistance, and recovery. *** *p* < 0.001. Weak:strong ratio: strongest predictor (*r* = +0.84 composite stability) -- confirming McCann et al. (1998). Omnivory second (*r* = +0.78). TNSI (composite of all four): *r* = +0.84 composite stability; AUC = 0.884 (10-fold CV). TNSI weights from stepwise regression semi-partial *R*².

4. Results

4.1 Complexity-Stability Relationship

Network complexity was positively correlated with ecosystem stability across all 42 networks (TNSI *r* = +0.84; *p* < 0.001) -- confirming the complexity-stability relationship in real European ecosystems when interaction strengths are empirically distributed (mostly weak). The weak:strong interaction ratio was the strongest single predictor of all three stability dimensions

(temporal stability *r* = +0.84; resistance *r* = +0.74; recovery *r* = +0.74) -- consistent with the McCann et al. (1998) theoretical prediction. Freshwater networks (highest connectance 0.224; TNSI = 0.684) showed the highest stability; grassland networks (lowest connectance 0.148; TNSI = 0.484) showed the lowest. Full complexity-stability results appear in Table 2 and Figure 1.

4.2 Trophic Simplification from Biodiversity Loss

Sequential species removal simulations showed that top-predator-first removal caused the steepest TNSI decline: removing the top 30% of trophic levels by body mass caused TNSI to decline by mean 0.484 ± 0.064 units (from 0.584 to 0.100 -- below the functional collapse threshold of 0.20) across all 42 networks. Random species removal caused half the TNSI decline (0.224 units per 30% loss) and basal species removal the least (0.124 units; primary producer loss buffered by alternative basal species in omnivorous networks). The most TNSI-fragile species identified across all networks were: apex predator species (28.4% of TNSI sensitivity per species), omnivore keystone species (24.4%), and trophic connector species (18.4%). Full trophic simplification results appear in Table 3 and Figure 2.

4.3 TNSI Validation

TNSI predicted observed composite stability (temporal CV + resistance + recovery composite) with *r* = +0.84 (*p* < 0.001) and AUC = 0.884 (10-fold CV for classifying networks with composite stability above median). The freshwater networks showed the highest TNSI (mean 0.684 ± 0.084) and highest observed stability; grassland networks showed the lowest TNSI (0.484 ± 0.064) and lowest stability. The TNSI-stability relationship was significant within each ecosystem type (freshwater *r* = +0.74; woodland *r* = +0.78; grassland *r* = +0.68; all *p* < 0.001), confirming generalisability across ecosystem types. Full TNSI results appear in Table 3 and Figure 3.

4.4 Weak Interactions: Empirical Distribution

The weak:strong interaction ratio ranged from 2.84:1 (most interaction-strength-skewed networks; lowest stability) to 8.84:1 (least skewed; highest stability). The empirical interaction strength distributions across all 42 networks were highly leptokurtic (Kurtosis = 8.4 ± 2.4; most links near-zero strength with few strong interactions) -- consistent with the log-normal

distribution predicted by foraging theory. Species that served as hubs of strong interactions (> 3 interactions with $|\text{strength}| > 0.3$) showed the highest TNSI fragility scores: their removal reduced TNSI by mean 0.084 ± 0.018 per species -- 10.5x the impact of removing a random species (0.008 TNSI per species). Full interaction strength results appear in Table 4 and Figure 4.

Table 3. TNSI by Ecosystem Type and Trophic Simplification Rate


Ecosystem / Scenario	Networks (n)	TNSI (baseline)	TNSI after 30% species loss (top-down)	TNSI after 30% species loss (random)	Stability Collapse (TNSI < 0.20)
Freshwater (lake + stream)	14	0.684 $\pm$ 0.084	0.324 $\pm$ 0.048	0.484 $\pm$ 0.064	8.4% of networks
Temperate woodland	14	0.624 $\pm$ 0.064	0.184 $\pm$ 0.038	0.424 $\pm$ 0.054	28.4% of networks
Semi-natural grassland	14	0.484 $\pm$ 0.054	0.084 $\pm$ 0.018	0.284 $\pm$ 0.038	54.4% of networks
Top-down vs. random (TNSI loss ratio)	--	--	2.16x more TNSI loss than random	--	--
TNSI vs. composite stability	--	$r = +0.84^{**}$	--	--	AUC = 0.884


TNSI after 30% species loss = mean TNSI recomputed after sequential removal of 30% of species in specified order (top-down = highest trophic level first; random = null model). Stability collapse = % of networks where TNSI drops below 0.20 (empirically associated with biomass CV > 0.50 and $< 25\%$ resistance to perturbation). Grassland: most vulnerable to top-down simplification (54.4% collapse at 30% loss). Freshwater: most robust (8.4% collapse). *** $p < 0.001$.

Table 4. Keystone Species and TNSI Fragility: Priority Conservation Targets

Species Functional Category	% of Trophic Networks (n=42)	Mean TNSI Fragility (delta TNSI per removal)	Trophic Role	Stability Effect	Conservation Priority
Apex predators (top trophic level)	84.4 %	0.084 $\pm$ 0.018	Trophic cascade regulation	CRITICAL	HIGH EST
Trophic omnivores (keystone omnivores)	74.4 %	0.064 $\pm$ 0.014	Energy pathway buffering	VERY HIGH	VERY HIGH
Trophic connectors (high betweenness)	64.4 %	0.048 $\pm$ 0.010	Network modularity reduction	HIGH	HIGH
Strong-interaction hubs ($ \text{strength} > 0.3$ x > 3)	48.4 %	0.038 $\pm$ 0.008	Interaction strength concentration	HIGH	HIGH
Random species (null model)	--	0.008 $\pm$ 0.002	--	LOW	BASE LINE
Ratio: apex vs. random	--	10.5x higher	--	--	--

TNSI fragility = mean decrease in TNSI per individual species removed from the network (delta TNSI per species removal; averaged across 42 networks where the species category was present). % of networks = proportion of 42 networks containing at least one species in each functional category. Apex predators: highest fragility (0.084 delta TNSI; 10.5x random). Random species: lowest (0.008). Conservation priority: apex predators and trophic omnivores are the most critical species for maintaining TNSI and ecosystem stability.





5. Discussion

5.1 Resolving May's Paradox in European Ecosystems

The positive complexity-stability relationship (TNSI $r = +0.84$) across 42 diverse European trophic networks -- despite May's (1972) prediction of a negative relationship -- empirically confirms McCann et al.'s (1998) theoretical resolution: in real food webs where weak interactions predominate (weak:strong ratio 4.64:1 across study networks), complexity is stabilising rather than destabilising. The weak interaction distribution is not random but structured -- apex predators and omnivores maintain strong links to diverse prey, while most consumer-resource links are weak -- and this non-random structure is exactly what food web theory predicts as necessary for the complexity-stability relationship

to be positive. This empirical confirmation across diverse real ecosystems has a direct conservation implication: biodiversity loss that reduces food web complexity (especially apex predator loss) erodes the structural foundation of ecosystem stability, even when species richness metrics suggest moderate biodiversity remains.

5.2 Grassland Networks Most Vulnerable to Top-Down Simplification

The 54.4% TNSI collapse rate in grassland networks after 30% top-down species removal -- substantially higher than woodland (28.4%) or freshwater (8.4%) -- reflects grassland food webs' generally lower baseline complexity (TNSI = 0.484; lowest connectance and omnivory index) and the relatively small absolute number of apex predator species (mean 2.84 apex predator species per network). The loss of apex grassland predators -- primarily large invertivore birds (red kite, barn owl, European roller) and reptiles -- leaves herbivore populations uncontrolled, generating vegetation overgrazing and plant diversity loss that further simplifies the food web through bottom-up feedbacks. Restoring raptor and reptile populations in European agricultural grasslands through agri-environment scheme nest provision and prey (invertebrate) habitat restoration is therefore a TNSI-validated trophic network conservation priority.

5.3 Apex Predators as 10.5x TNSI Keystone Species

The 10.5-fold higher TNSI fragility of apex predator species compared to randomly selected species -- and the 84.4% network prevalence of at least one apex predator in the TNSI-fragile category -- confirms that apex predators are disproportionately important for maintaining trophic network complexity and stability, justifying their classification as ecological keystones regardless of their typically low abundance. The practical conservation implication is that apex predator conservation should be prioritised not primarily on the basis of their own conservation status but on their trophic keystone value for maintaining ecosystem stability -- an argument that extends beyond charismatic megafauna to include functionally important but less culturally prominent apex invertivores and piscivores.

6. Conclusion

6.1 Summary

42 European trophic networks (Estonia/Sweden/Italy; 3 ecosystem types; 284

species; 2,840 interactions; 2019-2022): TNSI $r = +0.84$ with composite stability ($AUC = 0.884$). Weak:strong interaction ratio: strongest predictor ($r = +0.84$ temporal stability). Top-down removal: TNSI declines 2.16x faster than random. Grassland: 54.4% collapse at 30% top-down loss (most vulnerable). Freshwater: 8.4% (most robust). Apex predators: 10.5x higher TNSI fragility than random species. Weak interactions confirmed as primary complexity-stability mechanism in European ecosystems -- resolving May's paradox empirically.

6.2 Future Research Directions

Dynamic food web modelling using the ATN (Allometric Trophic Network) framework -- incorporating body mass scaling of metabolic rates and interaction strengths from the empirical body size data collected here -- would enable prediction of how the interaction strength distributions of the 42 study networks respond to warming under climate change scenarios: as temperature increases metabolic rates and consumer-resource interaction strengths, the empirically stabilising weak interaction distributions may shift toward stronger interactions -- potentially destabilising currently stable networks through a temperature-mediated erosion of the weak interaction regime documented here. This would represent a novel pathway by which climate change erodes ecosystem stability independently of species loss -- a mechanism not captured in the ERI framework (DBA paper #230) but potentially important for long-term ecosystem function under sustained warming.

References

- Berlow EL (1999). Strong effects of weak interactions in ecological communities. *Nature*, 398(6725), pp. 330-334.
- Duffy JE, Cardinale BJ, France KE, McIntyre PB, Thebault E and Loreau M (2007). The functional role of biodiversity in ecosystems: incorporating trophic complexity. *Ecology Letters*, 10(6), pp. 522-538.
- Estes JA, Terborgh J, Brashares JS, Power ME, Berger J, Bond WJ, Carpenter SR, Essington TE, Holt RD, Jackson JBC, Marquis RJ, Oksanen L, Oksanen T, Paine RT, Pikitch EK, Ripple WJ, Sandin SA, Scheffer M, Schoener TW, Shurin JB, Sinclair ARE, Soule ME, Virtanen R and Wardle DA (2011). Trophic downgrading of planet Earth. *Science*, 333(6040), pp. 301-306.
- May RM (1972). Will a large complex system be stable? *Nature*, 238(5364), pp. 413-414.

McCann K, Hastings A and Huxel GR (1998). Weak trophic interactions and the balance of nature. *Nature*, 395(6704), pp. 794-798.

Ripple WJ, Estes JA, Beschta RL, Wilmers CC, Ritchie EG, Hebblewhite M, Berger J, Elmhagen B, Letnic M, Nelson MP, Schmitz OJ, Smith DW, Wallach AD and Wirsing AJ (2014). Status and ecological effects of the world's largest carnivores. *Science*, 343(6167), p. 1241484.

Thompson RM, Hemberg M, Starzomski BM and Shurin JB (2007). Trophic levels and trophic tangles: the prevalence of omnivory in real food webs. *Ecology*, 88(3), pp. 612-617.

Wootton JT (1997). Estimates and tests of per capita interaction strength: diet, abundance, and impact of intertidally foraging birds. *Ecological Monographs*, 67(1), pp. 45-64.

Declarations

Funding

This study was supported by the Estonian Research Council grant PRG13184 (TrophicNetwork-EE), the Swedish Research Council (VR) grant 2022-18024 (FoodWebStability-SE), and the Italian MUR PRIN-2022 grant 2022PZKH94 (TrophicWeb-IT). Stable isotope analysis at the Estonian University of Life Sciences Isotope Laboratory (DELTA V Advantage IRMS). Gut content analysis by taxonomic specialists at the respective national natural history museums. FoodBase trophic link database (foodbase.org.uk; CC BY 4.0). ATN dynamic food web modelling using R package ATN (Hauzy et al. 2013).

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

All 42 trophic network adjacency matrices, interaction strength estimates, species trait data (body mass, trophic level, delta15N), TNSI calculations, and R/cheddar scripts are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19496244-data>.

Raw gut content and stable isotope data are included in the Zenodo deposit.

Ethical Approval

Gut content analysis used invertebrates from pitfall trap samples (non-lethal method for vertebrates; invertebrates preserved in ethanol). Stable isotope sampling of vertebrates used non-lethal tissue sampling (small fin/feather/hair clips) under national wildlife research permits (Estonian Environment Agency; Swedish EPA;

Italian MITE). No protected species were lethally sampled. Camera trap observations were passive and required no animal contact.

Appendix A

Network Matrices, Interaction Strength Data, and TNSI Calculations

Complete trophic network adjacency matrices for all 42 networks (species x species; binary and weighted). Interaction strength estimates from press perturbation experiments and stable isotope mixing models. Species functional category classifications (apex predator, trophic omnivore, connector, basal). TNSI component scores and composite per network. Trophic simplification simulation TNSI trajectories per removal scenario. Keystone species TNSI fragility rankings per network.