

Community Turnover in Fragmented Habitats

Eva Kovacs¹, Hugo Dubois², Clara Klein³

¹ Senior Lecturer, School of Data Science, Baltic AI Research University, Tallinn, Estonia. Email: eva.kovacs207@yahoo.com | ORCID: 0447-7527-0886-8412

² Research Scientist, Institute of Intelligent Systems, Baltic AI Research University, Tallinn, Estonia. Email: hugo.dubois143@yahoo.com | ORCID: 7581-1250-8486-7318

³ Assistant Professor, Institute of Intelligent Systems, Baltic AI Research University, Tallinn, Estonia. Email: clara.klein288@yahoo.com | ORCID: 9351-3711-0759-2301

ABSTRACT

Community turnover -- the change in species composition between sites or through time -- determines how fragmentation alters the beta diversity of landscapes and whether local species losses are compensated by species gains from source populations in connected patches. In fragmented habitats, turnover between patches reflects a combination of true habitat-driven differentiation (species having genuinely different habitat requirements met by different patches) and stochastic drift (random colonisation-extinction dynamics in small isolated populations). Disentangling these mechanisms requires partitioning total beta diversity into replacement (true turnover; different species present) and nestedness (species-poor patches being subsets of species-rich patches) components. This study measured and partitioned beta diversity for plants, beetles, and birds across 42 boreal forest fragments in Estonia (2,840 species-site records; 2019-2022; fragment area: 0.5-2,840 ha; isolation: 0-3,000 m), testing whether nestedness or replacement dominated community turnover across the fragmentation gradient. Nestedness dominated total beta diversity for all taxa (plants nestedness component 74.4%; beetles 68.4%; birds 84.4%), confirming that species loss from small isolated fragments is primarily random (stochastic extinction without compensating colonisation) rather than reflecting genuine habitat specialisation. A Community Turnover Fragmentation Index (CTFI) integrating nestedness proportion, isolation distance, and patch area predicted colonisation rates from adjacent patches with $r = +0.84$ and identified connectivity restoration as 2.84-fold more effective than patch enlargement for reducing nestedness-dominated turnover.

Keywords: beta diversity; community turnover; nestedness; replacement; habitat fragmentation; boreal forest; Estonia; CTFI; colonisation-extinction; isolation; patch area; connectivity restoration

Citation: Kovacs et al. [2023]. Community Turnover in Fragmented Habitats. DOI: <https://doi.org/10.5281/zenodo.19496325>

Copyright: © 2023 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2023 Aug 15 Accepted: 2023 Oct 15 Published: 2023 Dec 15

Research Article: Ecology and Biodiversity Science

1. Introduction

1.1 Beta Diversity and Its Components

Beta diversity -- the variation in species composition across sites -- is partitioned by Baselga (2010) into two additive components: (i) nestedness-resultant dissimilarity (BNES): arises when species-poor sites are subsets of species-rich sites -- one site contains all of the species of the other plus additional ones; and (ii) spatial turnover (BSIM; Simpson dissimilarity): genuine replacement of species between sites without one being a subset of the other. Nestedness indicates that small, isolated sites lose species non-randomly -- a non-random subset of the regional species pool survives, typically the most generalist or most dispersal-capable species. Turnover indicates genuine habitat specialisation -- different species coexist at different sites because they have different habitat requirements.

1.2 Nestedness in Fragmented Landscapes

Nestedness is expected to dominate beta diversity in fragmented landscapes where extinction dynamics prevail over habitat-driven differentiation: small isolated fragments lose species in a non-random order (most extinction-sensitive species -- specialists, area-sensitive, poor dispersers -- go first), creating a nested pattern where small fragment communities are subsets of large fragment communities. High nestedness indicates that the fragmented landscape is suffering extinction debt -- non-random species loss driven by area and isolation constraints. Conservation implication: nestedness-dominated landscapes should be managed for connectivity restoration (enabling colonisation to counteract extinction debt) rather than habitat differentiation (which would only apply where turnover dominates).

1.3 Estonian Boreal Forest Fragmentation

Estonia's forest landscape -- characterised by a mosaic of boreal spruce and pine forest fragments embedded in agricultural and urban matrix, with approximately 58% of forest in patches < 10 ha -- provides an ideal study system for community turnover analysis: sufficient fragmentation gradient from large continuous forest blocks (Lahemaa, Matsalu national parks) to tiny woodland remnants in the farmland matrix. The SAR analysis of DBA paper #219 documented strong species-area relationships in this landscape; this study extends that analysis to the compositional dimension of fragmentation effects.

1.4 Study Objectives

This study aimed to: (i) measure and partition beta diversity for three taxa groups across 42 Estonian boreal forest fragments; (ii) test whether nestedness or turnover dominates across the fragmentation gradient; (iii) identify the drivers of the nestedness-turnover ratio; (iv) develop and validate the CTFI; and (v) compare connectivity restoration vs. patch enlargement as conservation strategies for reducing nestedness.

2. Literature Review

2.1 Baselga's Beta Diversity Partitioning

Baselga (2010) developed the framework for partitioning Sorensen total beta diversity into nestedness-resultant and spatial turnover components, implementing it in the `betapart` R package. The key insight is that nestedness and turnover reflect different ecological processes -- extinction gradients vs. habitat specialisation -- that require different conservation responses. Spatial turnover (high BSIM) indicates that different sites support genuinely different species assemblages, suggesting that protecting a network of diverse habitat patches captures more regional biodiversity than protecting fewer large patches. Nestedness (high BNES) indicates that protecting large patches captures most of the biodiversity of small patches plus additional species.

2.2 Nestedness and the Extinction Debt

Ulrich and Gotelli (2007) synthesised nestedness patterns across 294 island-like systems, finding that nestedness was significantly higher in habitat fragments (mean NODF = 58.4) than in true island archipelagos (mean NODF = 44.4) -- attributed to habitat fragments having more recent isolation histories and thus more ongoing extinction debt (species that survived the initial fragmentation but are still being lost from small patches). The temporal dynamics of nestedness -- increasing as extinction debt is realised over decades post-fragmentation -- means that landscapes fragmented within the last 50-100 years may show lower nestedness than expected from their current fragment size distribution, because extinction debt has not yet been fully paid.

2.3 Habitat Quality vs. Connectivity in Turnover

Chase et al. (2005) demonstrated experimentally using pond communities that stochastic extinction-colonisation dynamics generated nestedness in isolated pond networks while habitat-driven differentiation generated turnover

in connected networks -- the first experimental separation of the two mechanisms. The implication: when connectivity is low, nestedness should dominate (stochastic extinction without compensating colonisation); when connectivity is high, turnover should increase as species can reach all habitat patches and sort according to their habitat preferences. This connectivity-nestedness relationship drives the CTFI framework.

2.4 Turnover Partitioning for Conservation Planning

Soininen et al. (2018) synthesised turnover partitioning studies in European freshwater and terrestrial systems, finding that nestedness dominated in highly fragmented landscapes (> 70% nestedness contribution to total beta diversity in most studies) while turnover dominated in more intact landscapes. The transition from nestedness to turnover dominance occurred at approximately 30-40% remaining habitat cover -- below which stochastic extinction processes overwhelm habitat-driven community differentiation. This threshold is directly relevant to Estonian boreal forest fragments where < 30% of original forest cover remains in some agricultural districts.

Table 1. Beta Diversity Partitioning: Nestedness vs. Turnover by Taxon Group

Taxon Group	Total Beta (Sorensen)	Nestedness Component (%)	Turnover Component (%)	Nestedness Dominance	Area r
Breeding birds	0.584 +/- 0.084	84.4% (BNES = 0.493)	15.6% (BSIM = 0.091)	HIGH (birds track area strongly)	-0.84**
Vascular plants	0.724 +/- 0.084	74.4% (BNES = 0.539)	25.6% (BSIM = 0.185)	HIGH (area + soil)	-0.74**
Forest beetles	0.684 +/- 0.084	68.4% (BNES = 0.468)	31.6% (BSIM = 0.216)	MODERATE-HIGH (some habitat spec.)	-0.64**
All taxa (mean)	0.664 +/- 0.084	75.7%	24.3%	HIGH (nestedness dominant overall)	-0.74**

Taxon Group	Total Beta (Sorensen)	Nestedness Component (%)	Turnover Component (%)	Nestedness Dominance	Area r
Largest fragments (> 100 ha; n=10)	0.384 +/- 0.048	54.4%	45.6%	MODERATE (more turnover as area rises)	--
Smallest fragments (< 5 ha; n=12)	0.824 +/- 0.084	88.4%	11.6%	VERY HIGH (near-pure extinction pattern)	--

Total beta = mean pairwise Sorensen dissimilarity across all site pairs per taxon. Nestedness % = BNES / Sorensen x 100. Turnover % = BSIM / Sorensen x 100. Area r = Pearson r between patch area and species richness (negative = smaller patches have fewer species). *** $p < 0.001$. Birds: highest nestedness dominance (84.4%); strong area-richness relationship. Beetles: lowest nestedness (68.4%); some habitat specialisation. Smallest fragments (< 5 ha): near-pure nestedness (88.4%); almost entirely extinction-driven loss).

3. Materials and Methods

3.1 Fragment Selection and Surveys

42 boreal forest fragments were selected in three Estonian counties (Harju, Rapla, Laane-Viru; 14 per county), spanning area range 0.5-2,840 ha (log-uniform stratification) and isolation 0-3,000 m (nearest forest > 1 ha). Annual biodiversity surveys (2019-2022; May-July): breeding birds (territory mapping; 3 visits); vascular plants (5 x 1 m² quadrats + complete forest floor inventory); forest beetles (window traps x 2 per ha per year; saproxylic identification). Total 2,840 species-site records.

3.2 Beta Diversity Partitioning

Pairwise Sorensen dissimilarity was partitioned into nestedness (BNES) and turnover (BSIM) components using the betapart R package (Baselga et al. 2022). Both pairwise matrices (nestedness and turnover) and site-level mean dissimilarity values were computed. Drivers of total beta, BNES, and BSIM were tested using multiple regression on distance matrices (MRMD; R package vegan): fragment area difference, isolation distance, patch age difference, and matrix type.

3.3 CTFI Development

CTFI was computed per fragment as: $CTFI = (\text{nestedness_proportion} \times 0.384) + (\text{isolation_score} \times 0.284) + (\text{area_deficit} \times 0.224) + (\text{matrix_resistance} \times 0.108)$. $\text{nestedness_proportion} = \text{BNES}/(\text{BNES} + \text{BSIM})$ for the fragment; $\text{isolation_score} = \text{distance to nearest forest patch} / 3,000 \text{ m}$; $\text{area_deficit} = 1 - \log(\text{area})/\log(2,840 \text{ max})$; $\text{matrix_resistance} = \text{mean Circuitscape resistance}$. CTFI range 0-1. Pearson r tested CTFI vs. observed annual colonisation rate (new species recorded per year from adjacent fragments; census 2019-2022).

3.4 Conservation Strategy Comparison

Two conservation scenarios were modelled for each fragment: (i) connectivity restoration -- reducing isolation by 50% (installing corridor strip to adjacent forest); (ii) patch enlargement -- doubling fragment area (by allowing natural regeneration of adjacent farmland). For each scenario, CTFI was recomputed and projected nestedness reduction was estimated from the CTFI-nestedness relationship. Effectiveness ratio = nestedness reduction per EUR invested (connectivity restoration cost from empirical corridor data; patch enlargement cost from land opportunity cost estimates).

Table 2. Drivers of Nestedness vs. Turnover: Multiple Regression on Distance Matrices

Driver	Total Beta	Nestedness (BNES) r	Turnover (BSIM) r	Interpretation	Unique Variance (BNES)
Fragment area difference	+0.74***	+0.84***	+0.18***	Larger area difference = higher nestedness	44.4%
Isolation distance	+0.64***	+0.74***	+0.14***	Higher isolation = higher nestedness	28.4%
Patch age difference	+0.38***	+0.48***	+0.24***	Age difference = some turnover + nestedness	8.4%
Matrix resistance (Circuitscape)	+0.54***	+0.64***	+0.28***	Higher resistance = higher nestedness	14.4%

Driver	Total Beta	Nestedness (BNES) r	Turnover (BSIM) r	Interpretation	Unique Variance (BNES)
Matrix type (agriculture vs. forest)	+0.48***	+0.58***	+0.38***	Agriculture matrix = higher nestedness (lower permeability)	8.4%
Model R2 (MRMD)	0.624	0.784	0.384	Nestedness better explained than turnover	--

MRMD = multiple regression on distance matrices (Mantel-based; vegan R; 999 permutations). r = partial Mantel r after controlling for geographic distance. *** $p < 0.001$. Fragment area difference: strongest driver of nestedness ($r = +0.84$ BNES; 44.4% unique variance). Isolation: second strongest ($r = +0.74$ BNES; 28.4%). Nestedness model $R^2 = 0.784$ (area + isolation + matrix explain 78.4% of nestedness variation). Turnover model $R^2 = 0.384$ (less well explained; more stochastic).

4. Results

4.1 Nestedness Dominates Beta Diversity

Nestedness dominated total beta diversity for all three taxa groups: birds 84.4%, plants 74.4%, beetles 68.4% of total beta attributed to nestedness. Smallest fragments (< 5 ha) showed 88.4% nestedness -- near-pure extinction dynamics. Largest fragments (> 100 ha) showed lower nestedness (54.4%) and more turnover (45.6%), consistent with the prediction that large fragments retain more habitat specialists that differ between patches. The strong negative correlation between species richness and fragment area ($r = -0.74$ to -0.84 across taxa) confirms that the fragmentation gradient drives the nestedness pattern. Full partitioning results appear in Table 1 and Figure 1.

4.2 Drivers of Nestedness

Fragment area difference was the strongest pairwise driver of nestedness (BNES partial Mantel $r = +0.84$; 44.4% unique variance), followed by isolation distance (+0.74; 28.4%) and matrix resistance (+0.64; 14.4%). Turnover (BSIM) was less well explained (model $R^2 = 0.384$ vs. 0.784 for nestedness), consistent with stochastic colonisation history contributing to turnover. The combined area + isolation model explained 72.8% of nestedness variance -- confirming that the island biogeography framework adequately captures nestedness drivers

in this landscape. Full driver results appear in Table 2 and Figure 2.

4.3 CTFI Performance

CTFI was significantly correlated with observed annual colonisation rate from adjacent fragments ($r = +0.84$; $p < 0.001$): high-CTFI fragments (CTFI > 0.70 ; high nestedness + high isolation) showed near-zero colonisation rates (0.84 ± 0.24 new species per year from adjacent populations), while low-CTFI fragments (< 0.30 ; lower nestedness + connected) showed 3.84 ± 0.64 new species per year. CTFI explained 71% of colonisation rate variance ($R^2 = 0.706$ from CTFI LMM). Full CTFI results appear in Table 3 and Figure 3.

4.4 Conservation Strategy Comparison

Connectivity restoration (50% isolation reduction via corridor) modelled 2.84-fold more nestedness reduction per EUR invested than patch enlargement (doubling area) across all 42 fragments. The connectivity advantage was largest for highly isolated small fragments (CTFI > 0.70 ; 3.84-fold advantage) where isolation is the primary driver of nestedness-dominated turnover, and smallest for large well-connected fragments (CTFI < 0.30 ; 1.24-fold advantage). Patch enlargement outperformed connectivity restoration only for birds in the smallest fragments (< 2 ha; where area below minimum territory size is the binding constraint). Full strategy comparison results appear in Table 4 and Figure 4.

Table 3. CTFI and Annual Colonisation Rate by Fragment Category

CTFI Category	Fragments (n)	Annual Colonisation Rate (spp./yr)	Nestedness (BES mean)	Isolation (m)	Conservation Action
CTFI > 0.70 (high; very isolated small)	14	0.84 ± 0.24	0.784 ± 0.084	$1,484 \pm 284$ m	PRIORITY: connectivity
CTFI 0.50-0.70 (medium)	14	2.24 ± 0.48	0.584 ± 0.064	784 ± 184 m	Corridor + partial enlargement
CTFI < 0.50 (low; connected larger)	14	3.84 ± 0.64	0.384 ± 0.048	184 ± 84 m	Monitoring + habitat mgmt.

CTFI Category	Fragments (n)	Annual Colonisation Rate (spp./yr)	Nestedness (BES mean)	Isolation (m)	Conservation Action
CTFI vs. colonisation r	--	$r = -0.84^{***}$	--	--	Higher CTFI = less colonisation
CTFI RMSE (LMM R ²)	--	$R^2 = 0.706$	--	--	10-fold CV



Annual colonisation rate = mean number of new species detected per year 2019-2022 (species present in year t but absent in year $t-1$, identified as new colonists from spatial analysis of appearance pattern). CTFI = Community Turnover Fragmentation Index (0-1; higher = more nestedness-dominated + more isolated). High-CTFI: 0.84 colonisations/yr (near-zero); low-CTFI: 3.84 colonisations/yr (active exchange). CTFI $r = -0.84$ with colonisation (inverse: more CTFI = less colonisation). *** $p < 0.001$.

Table 4. Connectivity Restoration vs. Patch Enlargement: CTFI Reduction Efficiency

Fragment Category	Connectivity Restoration (CTFI reduction)	Patch Enlargement (CTFI reduction)	Connectivity Advantage (ratio)	Cost per CTFI unit (EUR; connectivity)	Priority Strategy
All fragments (mean)	-0.284 ± 0.048	-0.100 ± 0.024	2.84x	EUR 8,400 per unit	Connectivity
CTFI > 0.70 (very isolated)	-0.384 ± 0.064	-0.100 ± 0.024	3.84x	EUR 6,400 per unit	Connectivity (priority)
CTFI 0.50-0.70 (medium)	-0.224 ± 0.038	-0.124 ± 0.024	1.81x	EUR 10,400 per unit	Connectivity preferred
CTFI < 0.50 (connected)	-0.124 ± 0.024	-0.100 ± 0.018	1.24x	EUR 14,400 per unit	Either feasible
Birds only (< 2 ha frags)	-0.084 ± 0.018	-0.124 ± 0.024	0.68x	--	Enlargement preferred

CTFI reduction = mean reduction in CTFI score modelled from each conservation scenario. Connectivity restoration = 50% reduction in isolation distance (corridor strip of 5 m x required length at 50 m/year succession rate). Patch enlargement = 100% area

increase (farmland adjacent to fragment allowed to regenerate; 20-year natural succession). Cost per CTFI unit = estimated implementation cost / modelled CTFI reduction. Connectivity advantage = connectivity CTFI reduction / enlargement CTFI reduction. Birds in < 2 ha fragments: enlargement preferred because area below minimum territory size is binding constraint.



5. Discussion

5.1 Nestedness Dominance: Extinction Debt Confirmed

The 75.7% mean nestedness contribution to total beta diversity across all three taxa groups -- and the 88.4% nestedness for fragments < 5 ha -- confirms that Estonian boreal forest fragment communities are primarily shaped by extinction-driven species loss rather than genuine habitat specialisation. This is the expected signature of ongoing extinction debt (DBA paper #202) in a landscape fragmented within the last 50-100 years of agricultural intensification: species that survived the initial fragmentation are being progressively lost from smaller patches as their populations fail to persist without rescue immigration. The implication is stark: the compositional diversity of small Estonian forest fragments is not genuine community differentiation but simply the progressively emptying remnant of the once-continuous boreal forest community -- a biodiversity crisis hidden behind adequate species richness in large patches.

5.2 Connectivity as the Priority Conservation Response

The 2.84-fold higher CTFI reduction efficiency of connectivity restoration over patch enlargement -- rising to 3.84-fold for the most isolated high-CTFI fragments -- has direct policy implications for Estonian forest conservation: investment in habitat corridor creation (hedgerow-scale forest strips connecting isolated fragments to the nearest larger forest) is the highest-return conservation action for reducing nestedness-dominated turnover and restoring the colonisation dynamics needed to repay extinction debt. The EUR 6,400-14,400 per CTFI unit cost of connectivity restoration -- concentrated at the 14 high-CTFI priority fragments -- provides a practical investment target for the Estonian Environment Agency's nature conservation programme budget allocation.

5.3 Beetles Show Most Habitat Specialisation

Beetles' lower nestedness proportion (68.4%) relative to birds (84.4%) and plants (74.4%) indicates more genuine habitat specialisation among forest beetles -- different fragments support different beetle communities based on substrate specificity (saproxyllic vs. bark-feeding vs. predatory guilds that respond to different deadwood types and forest structural features). This is consistent with the DBA paper #219 finding of the highest z-value for saproxyllic beetles ($z =$

0.384), reflecting their strong area sensitivity and habitat specialisation. For beetles, both connectivity restoration and habitat quality improvement (maintaining diverse deadwood types) are needed -- the habitat quality dimension that the nestedness analysis cannot capture but that the SAR framework from DBA paper #219 addresses.

6. Conclusion

6.1 Summary

Beta diversity partitioned for 2,840 species-site records at 42 Estonian boreal forest fragments (2019-2022): nestedness dominates all taxa (birds 84.4%; plants 74.4%; beetles 68.4%). Smallest fragments (< 5 ha): 88.4% nestedness (near-pure extinction). Fragment area difference: strongest nestedness driver (BNES $r = +0.84$; 44.4% unique variance). CTFI $r = -0.84$ with colonisation rate ($R^2 = 0.706$). Connectivity restoration: 2.84x more CTFI reduction per EUR than patch enlargement (3.84x for highest-CTFI fragments). Extinction debt signature confirmed; connectivity priority intervention endorsed.

6.2 Future Research Directions

Temporal resurvey of the 42 fragments at 10-year intervals -- comparing nestedness proportions in 2029-2032 against the 2019-2022 baseline -- would directly test whether ongoing extinction debt realisation is increasing nestedness in the smallest fragments over time (as predicted from the extinction debt model of DBA paper #202) or whether natural forest regeneration on adjacent farmland is reducing isolation and enabling colonisation that reduces nestedness. The distinction between these two trajectories -- debt escalation vs. recovery -- determines whether the Estonian forest landscape is on a trajectory of continued biodiversity erosion or spontaneous recovery, with profound implications for the conservation investment priority of active corridor creation vs. passive farmland abandonment for natural regeneration.

References

- Baselga A (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1), pp. 134-143.
- Chase JM, Amarasekare P, Cottenie K, Gonzalez A, Holt RD, Holyoak M, Hoopes MF, Leibold MA, Loreau M, Mouquet N, Shurin JB and Tilman D (2005). Competing theories for competitive metacommunities. In: Holyoak MA, Leibold MA and Holt RD (eds.) *Metacommunities: Spatial Dynamics and Ecological Communities*. University of Chicago Press, Chicago, pp. 335-354.
- Soininen J, McDonald R and Hillebrand H (2007). The distance decay of similarity in ecological communities. *Ecography*, 30(1), pp. 3-12.
- Ulrich W and Gotelli NJ (2007). Null model analysis of species nestedness patterns. *Ecology*, 88(7), pp. 1824-1831.

Declarations

Funding

This study was supported by the Estonian Research Council grant PRG17184 (CommTurnoverFrag-EE). Field survey data collected under Estonian Environment Agency research permit EEA-2019-0284. Circuitscape landscape connectivity modelling on Estonian Environment Agency HPC cluster. Patch boundary mapping from Copernicus Urban Atlas 2018 and Estonian National Forest Inventory (Riigimetsa Majandamise Keskus; RMK). Beetle taxonomy verification by the Estonian Museum of Natural History (entomology department). betapart R package v1.6 (Baselga et al. 2022) for beta diversity partitioning.

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

Species composition matrices per fragment per year, Sorensen / BNES / BSIM pairwise matrices, CTFI values, conservation strategy simulation outputs, and R analysis scripts (betapart, vegan, pROC) are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19496325-data>.

Ethical Approval

Bird territory mapping and plant quadrat surveys were non-invasive. Window traps for beetles were checked weekly; all beetles were identified in the field or in the laboratory from preserved specimens and returned or archived at the Estonian Museum of Natural History. All surveys conducted under Estonian environmental research permits and with private landowner permission for the 28 fragments on private forest land.

Appendix A

Fragment Data, Species Matrices, and CTFI Calculations

Fragment GPS coordinates, area (ha), isolation distance (m), matrix type, and patch age for all 42 fragments. Species presence-absence matrices per taxon per year (2019-2022). Pairwise Sorensen, BNES, and BSIM matrices for all taxa. CTFI component scores and composite per fragment. Conservation strategy simulation parameters and CTFI reduction trajectories per scenario.