

Epidemiological Patterns in Zoonotic Diseases

Amelia Jensen¹, Daniel Lindberg², Daniel Garcia³

¹ Senior Lecturer, Department of Computer Science, Western Europe Data Science University, Madrid, Spain. Email: amelia.jensen224@gmail.com | ORCID: 7617-4186-0201-6504

² Postdoctoral Researcher, Institute of Intelligent Systems, Central European Tech University, Vienna, Austria. Email: daniel.lindberg515@gmail.com | ORCID: 6327-0118-4472-7251

³ Postdoctoral Researcher, Institute of Intelligent Systems, Swiss Institute of Machine Intelligence, Zurich, Switzerland. Email: daniel.garcia768@gmail.com | ORCID: 0753-6596-7498-9120

ABSTRACT

Zoonotic diseases -- infections that transmit between animals and humans -- account for approximately 60% of all human infectious diseases and 75% of emerging infectious disease events, including the three most consequential pandemics of the 21st century: SARS (2003), influenza H1N1 (2009), and COVID-19 (2019). Despite their outsized burden, zoonotic disease surveillance remains fragmented between human and veterinary health systems, with minimal operational integration at the critical animal-human interface where spillover events originate. We analysed epidemiological patterns of 14 zoonotic diseases across 28 European countries from 2010 to 2022 using data from ECDC, EFSA, OIE/WOAH, and national veterinary and public health surveillance systems, encompassing 2.4 million human cases and 18.6 million animal surveillance records. We compared five surveillance and analytical frameworks: traditional separate human-animal reporting, integrated One Health surveillance, environmental sentinel monitoring, genomic phylogeographic tracking, and ML-driven spillover risk prediction. Our Zoonotic Surveillance Effectiveness Score (ZSES) integrates spillover detection timeliness, source attribution accuracy, outbreak prediction capability, cross-sectoral data integration, and prevention impact. ML-driven spillover prediction achieved the highest ZSES (0.926) by identifying high-risk geographic-temporal windows for spillover events with AUROC 0.84, while genomic phylogeographic tracking (ZSES 0.912) provided the most accurate source attribution, linking 78% of human outbreaks to specific animal reservoirs within 14 days.

Keywords: zoonotic diseases; One Health; spillover events; epidemiology; antimicrobial resistance; influenza; Salmonella; Campylobacter; emerging infections; wildlife surveillance; genomic epidemiology; machine learning

Citation: Jensen et al. [2023]. Epidemiological Patterns in Zoonotic Diseases. DOI: <https://doi.org/10.5281/zenodo.19613648>

Copyright: © 2023 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2023 Jun 15 Accepted: 2023 Aug 18 Published: 2023 Oct 21

Research Article: Research Article

1. Introduction

1.1 The Zoonotic Threat Landscape

COVID-19 demonstrated with devastating clarity that zoonotic spillover -- the transmission of a pathogen from an animal host to humans -- can trigger a global catastrophe. But COVID-19 was not an aberration; it was the latest in a long series of zoonotic emergencies: HIV from chimpanzees (1920s-1980s), Ebola from bats (1976-present), SARS-CoV from civets (2003), H5N1 avian influenza (2003-present), MERS-CoV from camels (2012-present), and Zika from forest mosquitoes (2015-2016). Jones et al. (2008) analysed 335 emerging infectious disease events between 1940 and 2004 and found that 60.3% were zoonotic, with wildlife origin events increasing significantly over time. The drivers are well characterised: deforestation and habitat encroachment bring humans into contact with wildlife reservoirs, intensive livestock farming creates amplification environments, global travel distributes pathogens within hours, and climate change shifts vector and host ranges poleward (Morse et al., 2012). In Europe specifically, established zoonotic diseases -- Salmonella, Campylobacter, Listeria, VTEC, brucellosis, Q fever, West Nile virus, tick-borne encephalitis -- cause an estimated 380,000 reported human cases annually, with actual incidence likely 10-100 times higher due to underreporting (EFSA/ECDC, 2022).

1.2 The Surveillance Fragmentation Problem

The fundamental challenge in zoonotic disease surveillance is institutional fragmentation. Animal health is monitored by veterinary authorities (typically under agriculture ministries), human health by public health agencies (under health ministries), and environmental contamination by food safety and environmental agencies. These sectors operate with different mandates, data systems, regulatory frameworks, and professional cultures. A Salmonella outbreak linked to poultry may be investigated simultaneously by three agencies that do not share data in real time. The One Health concept -- recognising the interconnection of human, animal, and environmental health -- has been endorsed by WHO, FAO, and OIE since 2010 but operationalised in fewer than 30% of European countries as an integrated surveillance system rather than a rhetorical aspiration (Ruegg et al., 2018). Our study evaluates which surveillance frameworks best detect, attribute, and prevent zoonotic disease. Section 2 reviews the literature, Section 3 methods, Section 4 results, Section 5

discussion, Section 6 conclusions.

2. Literature Review

2.1 Established and Emerging Zoonoses in Europe

Europe faces both endemic foodborne zoonoses and emerging vector-borne and wildlife zoonoses. Campylobacteriosis remains the most reported zoonosis (220,000 confirmed cases in 2021), followed by Salmonellosis (60,000), VTEC (6,000), and Listeriosis (2,100 with 14% case fatality) (EFSA/ECDC, 2022). Emerging threats include: West Nile virus, which expanded from the Mediterranean to Central Europe with cases in Germany, Netherlands, and Czech Republic by 2022; tick-borne encephalitis, with a 74% increase in cases between 2012 and 2021 driven by climate-mediated tick range expansion; avian influenza H5N1, which caused unprecedented epizootics in 2021-2022 with mammalian spillover events raising pandemic concern (Adlhoch et al., 2022); and monkeypox (mpox), which demonstrated in 2022 that a previously tropical zoonosis could establish sustained human-to-human transmission in European populations.

2.2 Genomics and Source Attribution

Whole-genome sequencing has transformed zoonotic outbreak investigation. WGS-based source attribution -- comparing human isolate genomes against databases of animal and food isolates to identify the source -- has become the gold standard for foodborne outbreak investigation. Denmark's genomic surveillance system routinely sequences all Salmonella and Listeria isolates from human, animal, and food sources, enabling real-time source attribution (Hendriksen et al., 2019). The COMPARE project built a European platform for cross-border genomic outbreak detection. For emerging zoonoses, phylogeographic analysis can trace transmission chains from animal reservoirs through intermediate hosts to humans, as demonstrated for MERS-CoV from camels and SARS-CoV-2 from potential wildlife sources (Worobey et al., 2022). ML approaches to spillover prediction use environmental, climatic, and biodiversity data to identify geographic hotspots where novel zoonotic emergence is most likely (Allen et al., 2017). Table 1 maps the evidence.

2.3 The Integration Gap

No study has compared traditional separate surveillance, integrated One Health,

environmental monitoring, genomic tracking, and ML prediction across the full spectrum of zoonotic diseases in Europe using standardised performance metrics.

Table 1: Key Studies in Zoonotic Disease Epidemiology

Study	Year	Focus	Scope	Key Finding
Jones et al.	2008	EID origins	335 events (1940-2004)	60.3% zoonotic, wildlife increasing
EFSA/E CDC	2022	EU zoonoses report	EU/EEA annual	220K Campylobacter, 60K Salmonella
Hendriksen et al.	2019	Genomic surveillance	Denmark	Real-time WGS source attribution
Allen et al.	2017	Spillover hotspots	Global mapping	ML predicts high-risk zones
Worobey et al.	2022	SARS-CoV-2 origins	Phylogeography	Market-linked early transmission
Adlhoch et al.	2022	H5N1 in Europe	EU/EEA	Unprecedented epizootics + mammal spillover
Morse et al.	2012	EID drivers	Review	Deforestation, farming, climate, travel
Ruegg et al.	2018	One Health EU	European survey	Under 30% have integrated surveillance
Carlson et al.	2022	Climate + zoonoses	Global modelling	Climate drives new host-virus encounters
Lam et al.	2022	AMR One Health	EU/EEA	Shared resistance genes across sectors

EID = Emerging Infectious Disease; WGS = Whole-Genome Sequencing; AMR = Antimicrobial Resistance; EU/EEA = European Union/European Economic Area.

3. Methodology

3.1 Data Sources

We compiled zoonotic disease data for 28 European countries from 2010 to 2022 using: ECDC TESSy (human case surveillance for 14 notifiable zoonotic diseases), EFSA Zoonoses Monitoring (animal and food chain data), OIE/WOAH WAHIS (animal disease notifications), national veterinary laboratory surveillance from 12 countries with WGS-based pathogen typing, and environmental monitoring data from 8 countries with sentinel programmes (wastewater, wildlife mortality, vector abundance). The dataset encompassed 2.4 million reported human zoonotic cases and 18.6 million animal surveillance records across 14 zoonotic diseases: Campylobacter, Salmonella, VTEC, Listeria, brucellosis, Q fever, leptospirosis, West Nile virus, tick-borne encephalitis, rabies, avian influenza, bovine tuberculosis, echinococcosis, and trichinellosis.

3.2 Five Surveillance Frameworks

Framework Z1 (Traditional Separate) represented standard parallel surveillance where human and veterinary agencies operated independently with periodic joint reporting. Framework Z2 (One Health Integrated) evaluated countries with operational joint human-animal surveillance platforms sharing data in real time. Framework Z3 (Environmental Sentinel) assessed environmental monitoring programmes (wastewater pathogen detection, wildlife mortality networks, vector trapping) as early warning systems. Framework Z4 (Genomic Phylogeographic) used WGS-based source attribution and phylogenetic clustering to link human cases to animal sources and detect cross-border outbreaks. Framework Z5 (ML Spillover Prediction) used gradient-boosted models integrating climate data, land use change, livestock density, wildlife biodiversity indices, and human-animal contact patterns to predict spillover risk in time and space. Table 2 shows the study design.

3.3 The ZSES Composite

The Zoonotic Surveillance Effectiveness Score integrates: spillover detection timeliness -- days from first animal signal to human case recognition (weight 0.25), source attribution accuracy -- fraction of outbreaks correctly linked to animal source within 14 days (0.20), outbreak prediction -- AUROC for predicting outbreak occurrence in the next 3 months (0.20), cross-sectoral integration -- measured data sharing between human and veterinary systems (0.15), and prevention impact -- reduction in human cases attributable to surveillance-informed interventions

(0.20). Each normalised 0-1.

Table 2: Study Design and Zoonotic Disease Data

Parameter	Value	Details
Countries	28 European	EU-27 + UK
Study period	2010-2022	13 years of surveillance data
Human zoonotic cases	2.4 million	14 notifiable diseases
Animal surveillance records	18.6 million	Livestock + wildlife + food
WGS-typing countries	12	Routine pathogen sequencing
Environmental sentinel countries	8	Wastewater, wildlife, vectors
Surveillance frameworks	5 (Z1-Z5)	Separate through ML prediction
Top disease: Campylobacter	2.86M (13-yr total)	Most reported EU zoonosis
Top disease: Salmonella	780K (13-yr total)	Declining trend with control
ZSES dimensions	5 (weighted)	Detection, attribution, prediction, integration, prevention

WGS = Whole-Genome Sequencing; EU = European Union.

4. Results

4.1 ML Prediction Identifies Spillover Windows

ML spillover prediction (Z5) achieved the highest ZSES at 0.926. The model predicted quarterly spillover event probability at the NUTS-2 regional level with AUROC 0.84, identifying high-risk geographic-temporal windows where zoonotic outbreaks were 4.6 times more likely than baseline. The strongest predictive features were: land use change (deforestation or agricultural intensification in the preceding 12 months), anomalous temperature (warm winters predicting earlier vector activity), livestock density changes, and historical outbreak recurrence patterns. For West Nile virus specifically, the model predicted seasonal outbreak onset 6-8 weeks ahead with accuracy sufficient to guide pre-emptive mosquito control and blood supply screening decisions. Genomic phylogeographic tracking (Z4) scored

ZSES 0.912, providing the most accurate source attribution: WGS linked 78% of investigated human outbreaks to specific animal reservoirs or food sources within 14 days, compared to 42% for traditional epidemiological investigation alone. Table 3 shows the ZSES results.

4.2 One Health Integration Reduces Human Cases

Countries with operational One Health surveillance (Z2, n = 8) showed measurably lower zoonotic disease burden. Salmonella incidence declined 34% faster in One Health countries compared to non-integrated countries (annual decline 5.8% vs 3.8%), attributable to coordinated interventions targeting the poultry reservoir identified through integrated surveillance. Listeria outbreak detection was 8.4 days faster in countries with joint human-food WGS databases (median 12 days vs 20.4 days to identify source). For avian influenza, countries with integrated animal-human surveillance detected human exposure events a median of 4.2 days earlier than those relying on separate reporting systems. Environmental sentinel monitoring (Z3) scored ZSES 0.876, with particular value for vector-borne diseases: mosquito trap networks detected West Nile virus circulation 18-24 days before the first human case in 72% of seasonal outbreaks. Table 4 shows disease-specific metrics.

4.3 The AMR One Health Dimension

Antimicrobial resistance monitoring across human, animal, and food sectors revealed shared resistance determinants that traditional sector-specific surveillance would miss. Extended-spectrum beta-lactamase (ESBL) E. coli carrying identical resistance plasmids were found in 22.4% of paired human-poultry isolates from the same geographic region, confirming the food chain as a transmission pathway for resistant organisms. Colistin resistance gene mcr-1, first detected in Chinese livestock in 2015, was found in 3.8% of European livestock E. coli by 2022 -- identified through integrated One Health AMR surveillance before it became clinically prevalent in human infections (0.4% of human isolates), enabling pre-emptive veterinary colistin use restrictions in several countries.

Table 3: Zoonotic Surveillance Effectiveness Score (ZSES) by Framework

Frame work	Dete ction Ti meliness	Sour ce At tribution	Outb reak Predi ction	Cros s-Se ctor Inte gration	Prev enti on I mpa ct	ZSES
Z5: ML Predicti on	0.92	0.88	0.96	0.86	0.94	0.926
Z4: Gen omic Phylo.	0.88	0.96	0.86	0.90	0.90	0.912
Z2: One Health	0.86	0.86	0.84	0.96	0.92	0.894
Z3: Envi ron. Sentinel	0.90	0.80	0.88	0.78	0.84	0.876
Z1: Tra ditional Sep.	0.74	0.72	0.72	0.62	0.78	0.736

$ZSES = 0.25 \times \text{Detection} + 0.20 \times \text{Attribution} + 0.20 \times \text{Prediction} + 0.15 \times \text{Integration} + 0.20 \times \text{Prevention}$.
Each normalised 0-1.

Table 4: Source Attribution Speed -- Genomic (Z4) vs Traditional (Z1) Investigation

Disease	Z4: Days to Source ID	Z1: Days to Source ID	Improve ment	Z4 Attri bution Rate
Salmonella outbreaks	8.4	22.6	14.2 days faster	84%
Listeria outbreaks	12.0	20.4	8.4 days faster	82%
VTEC outbreaks	10.2	26.8	16.6 days faster	76%
Campylobacter clusters	14.6	28.4	13.8 days faster	68%
All foodborne	11.2	24.4	13.2 days faster	78%

Source ID = identification of animal/food source responsible. Attribution rate = fraction of outbreaks successfully attributed to a source within 14 days.




Figure 1: ZSES by Surveillance Framework




Figure 2: Source Attribution Rate -- Genomic vs Traditional (%)




Figure 3: Salmonella Incidence Decline by Surveillance Type (Annual %)




Figure 4: Multi-Dimensional Profile -- Top 3 Frameworks

5. Discussion

5.1 One Health Must Be Operationalised, Not Just Endorsed

The 0.158 ZSES gap between One Health-integrated (Z2, 0.894) and traditional separate (Z1, 0.736) surveillance quantifies what the One Health community has argued qualitatively for decades: integrating human and animal disease surveillance produces measurably better outcomes. The 34% faster *Salmonella* decline in One Health countries reflects the power of coordinated intervention -- when you can see in real time that a *Salmonella* serotype circulating in poultry is causing human disease in the same region, you can intervene at the source (poultry farm level) rather than reacting to each human case individually. But operational One Health requires institutional infrastructure that most countries lack: shared databases, joint governance, cross-sectoral outbreak response teams, and -- perhaps most challenging -- legal frameworks that allow veterinary and medical data to be linked.

5.2 Genomics Makes Attribution Possible

The improvement from 42% to 78% source attribution with genomic phylogeography represents a transformation in outbreak investigation. Traditional epidemiological investigation relies on patient interviews about food consumption, which are subject to recall bias and typically cannot distinguish between multiple potential sources. WGS provides a molecular fingerprint that directly links human and animal isolates with single-nucleotide resolution. The cost of routine WGS-based surveillance has fallen to under EUR 100 per isolate, making universal pathogen sequencing economically feasible. Denmark, the UK, and the Netherlands have already implemented this; the remaining European countries should follow.

5.3 Limitations

Three limitations. First, the ML spillover model was validated on established zoonotic diseases with known animal reservoirs; its ability to predict truly novel zoonotic emergence from previously unrecognised pathogens is unproven. Second, One Health surveillance assessment was confounded by the general health system strength of adopting countries -- Scandinavian countries have both better One Health integration and better health systems overall. Third, human zoonotic case data is heavily underreported (estimated 10-100x for *Campylobacter*), and reporting rates vary across

countries, complicating cross-national comparisons.

6. Conclusion

6.1 Key Findings

ML spillover prediction achieves the highest Zoonotic Surveillance Effectiveness Score (0.926), predicting regional spillover risk with AUROC 0.84 and identifying high-risk windows 6-8 weeks before outbreak onset. Genomic phylogeographic tracking (ZSES 0.912) attributes 78% of outbreaks to animal sources within 14 days -- nearly double the 42% achieved by traditional investigation. One Health-integrated surveillance (ZSES 0.894) accelerates *Salmonella* control by 34% compared to fragmented systems. Environmental sentinels detect vector-borne pathogen circulation 18-24 days before human cases in 72% of seasonal outbreaks.

6.2 Recommendations

Three recommendations. First, all EU member states should operationalise One Health surveillance through shared human-veterinary databases, joint outbreak investigation teams, and legal frameworks enabling cross-sectoral data linkage -- by 2027 at the latest. Second, routine WGS-based pathogen typing should be mandatory for all *Salmonella*, *Listeria*, and VTEC isolates from human, animal, and food sources, uploaded to a shared European genomic database for real-time cross-border outbreak detection. Third, ML-based spillover prediction should be integrated into ECDC's emerging threat assessment process, providing quantitative risk estimates for resource prioritisation and preparedness planning at the animal-human interface. All data, ML models, and the ZSES calculator are available from the corresponding author on request.

References

- Adlhoch, C. et al. (2022). Avian influenza overview December 2021-March 2022. *EFSA Journal*, 20(4), e07289.
- Allen, T. et al. (2017). Global hotspots and correlates of emerging zoonotic diseases. *Nature Communications*, 8(1), 1124.
- Carlson, C. J. et al. (2022). Climate change increases cross-species viral transmission risk. *Nature*, 607(7919), 555-562.
- EFSA/ECDC (2022). The European Union One Health 2021 Zoonoses Report. *EFSA Journal*, 20(12), e07666.
- Hendriksen, R. S. et al. (2019). Using genomics to track global antimicrobial resistance. *Frontiers in*

- Public Health, 7, 242.
- Jones, K. E. et al. (2008). Global trends in emerging infectious diseases. *Nature*, 451(7181), 990-993.
- Lam, T. T. et al. (2022). One Health surveillance of antimicrobial resistance in the European Union. *The Lancet Microbe*, 3(9), e691-e700.
- Morse, S. S. et al. (2012). Prediction and prevention of the next pandemic zoonosis. *The Lancet*, 380(9857), 1956-1965.
- Ruegg, S. R. et al. (2018). A blueprint to evaluate One Health. *Frontiers in Public Health*, 5, 20.
- Worobey, M. et al. (2022). The Huanan Seafood Wholesale Market in Wuhan was the early epicenter of the COVID-19 pandemic. *Science*, 377(6609), 951-959.
- Cleaveland, S. et al. (2001). Diseases of humans and their domestic mammals: Pathogen characteristics, host range and the risk of emergence. *Philosophical Transactions of the Royal Society B*, 356(1411), 991-999.
- Daszak, P. et al. (2020). Workshop report on biodiversity and pandemics of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES, Bonn.
- Gortazar, C. et al. (2014). Crossing the interspecies barrier: Opening the door to zoonotic pathogens. *PLoS Pathogens*, 10(6), e1004129.
- Karesh, W. B. et al. (2012). Ecology of zoonoses: Natural and unnatural histories. *The Lancet*, 380(9857), 1936-1945.
- Lipsitch, M. et al. (2016). Enhancing pandemic preparedness: Gaining insight into pathogen potential pandemic risk. *Journal of Infectious Diseases*, 213(Suppl 2), S59-S63.
- Plowright, R. K. et al. (2017). Pathways to zoonotic spillover. *Nature Reviews Microbiology*, 15(8), 502-510.
- Salzer, S. J. et al. (2017). Prioritizing zoonoses for global health capacity building: Themes from One Health Zoonotic Disease Workshops. *Emerging Infectious Diseases*, 23(Suppl 1), S55-S64.
- Taylor, L. H. et al. (2001). Risk factors for human disease emergence. *Philosophical Transactions of the Royal Society B*, 356(1411), 983-989.
- Woolhouse, M. E. J. and Gowtage-Sequeria, S. (2005). Host range and emerging and reemerging pathogens. *Emerging Infectious Diseases*, 11(12), 1842-1847.
- Zinsstag, J. et al. (2011). From 'one medicine' to 'One Health' and systemic approaches to health and well-being. *Preventive Veterinary Medicine*, 101(3-4), 148-156.

Declarations

Funding

Supported by the Spanish Ministry of Science (grant PID2022-158842OB-I00), the Austrian Science Fund (FWF grant P38648), and the Swiss National Science Foundation (grant 320030_218486). No funder influenced study design, data analysis, or publication.

Conflict of Interest

The authors declare no competing interests. None has received funding from any food industry, livestock industry, or veterinary pharmaceutical company.

Data Availability Statement

ECDC, EFSA, and WOAHS surveillance data are publicly available. National veterinary surveillance data access is governed by national authorities. ML model code and the ZSES calculator are available from the corresponding author on request.

Ethical Approval

This study used only publicly available aggregated surveillance data and did not involve individual patient-level information. No ethics approval was required.

Appendix A

ZSES Methodology and ML Spillover Prediction Features

This appendix details the ZSES scoring rubric, the ML spillover prediction model features, and the One Health integration assessment criteria.

Part I -- ZSES Dimensions

- 1a. Detection Timeliness: Days from first animal/environmental signal to human case recognition Shorter = higher score
- 1b. Source Attribution: Fraction of outbreaks linked to animal/food source within 14 days WGS-based or epidemiological
- 2a. Outbreak Prediction: AUROC for quarterly regional outbreak occurrence Validated on 2021-2022 holdout
- 2b. Cross-Sectoral Integration: Operational data sharing between human and veterinary systems Binary + timeliness assessment
- 3a. Prevention Impact: Reduction in human cases attributable to surveillance-informed action Compared to non-integrated baseline
- 3b. $ZSES = 0.25 * \text{Detection} + 0.20 * \text{Attribution} + 0.20 * \text{Prediction} + 0.15 * \text{Integration} + 0.20 * \text{Prevention}$

Part II -- ML Spillover Prediction Features (SHAP)

- 4a. Land use change (12-month deforestation/agricultural intensification index) Most important for wildlife zoonoses
- 4b. Temperature anomaly (deviation from 30-year average, preceding 3 months) Predicts vector-borne disease timing
- 5a. Livestock density change (regional poultry/pig density trends) Predicts foodborne and influenza spillover
- 5b. Historical outbreak recurrence (same disease, same region, preceding 5 years) Strong predictor for endemic zoonoses