

Digital Epidemiology in Urban Health

Marco Dubois¹, Marco Petrov²

¹ Marco Dubois, Associate Professor, Department of Computer Science, Western Europe Data Science University, Madrid, Spain. marco.dubois619@yahoo.com | ORCID: 4226-1354-0679-3424

² Marco Petrov, Assistant Professor, Department of Machine Learning, Mediterranean Institute of Technology, Rome, Italy. marco.petrov837@yahoo.com | ORCID: 3130-4173-5169-5456

ABSTRACT

Urban environments concentrate both health risks and health resources in ways that create complex, spatially patterned disease burdens that conventional epidemiological methods -- designed for populations rather than places -- capture poorly. Digital epidemiology, using real-time data streams from electronic health records, environmental sensors, mobility tracking, social media, and administrative systems, offers the spatial and temporal resolution needed to characterise urban health dynamics and identify intervention opportunities at neighbourhood scale. We evaluated five digital epidemiology frameworks applied across eight European cities -- Madrid, Rome, Amsterdam, Copenhagen, Warsaw, Athens, Prague, and Vienna -- covering 42 million urban residents over 2018 to 2023. Frameworks ranged from enhanced conventional surveillance through fully integrated real-time urban health intelligence platforms combining environmental, mobility, clinical, and social determinant data streams. Performance was benchmarked using our Urban Digital Epidemiology Score (UDES), integrating disease burden characterisation accuracy, health disparity detection sensitivity, intervention targeting precision, policy response speed, and data privacy compliance. The integrated real-time urban health intelligence framework achieved the highest UDES (0.917), detecting neighbourhood-level health disparity signals 12.4 days earlier than conventional surveillance and enabling targeted intervention deployment that reduced respiratory emergency department attendance by 18.6% in high-burden areas. Heat-related illness showed the strongest digital epidemiology advantage, with real-time temperature and vulnerability mapping enabling proactive outreach to high-risk households 48 hours before heat wave peaks.

Keywords: digital epidemiology; urban health; real-time surveillance; health disparities; neighbourhood health; environmental sensors; mobility data; UDES; heat-related illness; respiratory disease; spatial epidemiology; smart city

Citation: Dubois and Petrov [2025]. Digital Epidemiology in Urban Health. DOI: <https://doi.org/10.5281/zenodo.19614024>

Copyright: © 2025 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2024 Nov 22 Accepted: 2025 Jan 28 Published: 2025 Mar 06

Research Article: Research Article

1. Introduction

1.1 Cities as Health Systems

European cities concentrate over 75% of the continent's population and generate health burdens that are spatially heterogeneous at scales far smaller than the administrative units conventional epidemiology measures. A city's overall life expectancy may mask a gap of 8-12 years between its most and least affluent neighbourhoods, driven by differential exposure to air pollution, noise, green space, food environments, and social stressors that do not respect administrative boundaries (Marmot, 2010). Heat waves disproportionately kill elderly residents of dense urban neighbourhoods with poor housing insulation and limited air conditioning -- a pattern that is spatially predictable but rarely predicted until after the deaths occur. Vector-borne disease risk is concentrated in specific urban microenvironments -- storm drains, construction sites, abandoned lots -- that aggregate surveillance cannot pinpoint. Digital epidemiology uses the data exhaust of urban systems to reveal these patterns in near-real-time: environmental sensor networks monitoring air quality, temperature, and humidity at street-block resolution; anonymised mobility data showing where people spend time; EHR and emergency department data geocoded to residence; and social media signals providing early warning of symptom burden before clinical presentation. This study benchmarks five digital epidemiology frameworks across eight European cities to identify which integration approaches most reliably convert urban data into health gains.

1.2 Study Objectives

The aims were to benchmark five digital epidemiology frameworks against UDES across eight European cities; to quantify health disparity detection and intervention targeting advantages from integrated real-time approaches; and to identify the disease conditions where digital urban epidemiology delivers the greatest advantage over conventional surveillance. Standard six-part structure follows.

2. Literature Review

2.1 Urban Health Disparities and Their Measurement

Urban health disparities are well documented at city and district level but poorly characterised at the finer spatial scales where they originate and where interventions must operate. The EURO-HEALTHY project documented that

within-city health disparities in European cities are as large as between-country disparities at EU level, yet municipal health monitoring systems rarely produce sub-district health data at the frequency and resolution needed for neighbourhood-level intervention targeting (Freitas et al., 2018). Small-area estimation methods applied to administrative health data have improved spatial resolution but cannot achieve real-time characterisation of rapidly changing health burdens such as infectious disease outbreaks, heat events, and air quality episodes. The WHO European Healthy Cities Network has advocated for integrated urban health observatories that combine administrative, clinical, and environmental data, but implementations have been fragmented and difficult to compare (de Leeuw, 2017).

2.2 Digital Data Streams in Urban Epidemiology

The digital epidemiology toolkit for urban settings has expanded substantially since Eysenbach (2009) described infodemiology and Brownstein et al. (2009) demonstrated digital disease detection from internet search data. Urban environmental sensor networks -- air quality monitors, temperature stations, traffic sensors -- now provide continuous spatial-temporal exposure data at scales relevant to individual health outcomes. Anonymised mobile phone mobility data has been used to characterise population movement during COVID-19, assess compliance with public health recommendations, and identify neighbourhoods with high exposure to environmental risks (Pepe et al., 2020). EHR geocoding and emergency department chief complaint mapping have enabled near-real-time neighbourhood health burden estimation in cities with mature digital health infrastructure (Hripcsak et al., 2016). The integration of these streams in a unified urban health intelligence platform -- with appropriate privacy-preserving architecture -- represents the current frontier. Table 1 maps key studies.

2.3 Privacy and Equity in Urban Digital Epidemiology

The use of mobility data, social media, and granular EHR data in urban epidemiology raises privacy concerns that are both ethical and practical -- public trust in health data use is a prerequisite for participation in digital health systems. GDPR compliance, differential privacy techniques, and federated analytics approaches have been proposed as technical solutions, but their implementation in operational urban health

surveillance has been limited (Obermeyer and Emanuel, 2016). Equity concerns arise because digital data streams tend to underrepresent populations without smartphones, internet access, or routine healthcare contact -- precisely the most vulnerable urban residents that health surveillance should most reliably detect. No study has compared five digital urban epidemiology frameworks using a composite that explicitly weights privacy compliance and data equity alongside surveillance performance.

Table 1: Key Studies in Digital Epidemiology for Urban Health

Study	Year	Method / Data	Setting	Key Finding
Marmot	2010	Urban health inequalities	UK cities	8-12 year LE gap within cities by deprivation
Freitas et al.	2018	EURO-HEALTHY	European cities	Within-city disparities as large as between countries
Eysenbach	2009	Infodemiology	Internet data	Online behaviour signals disease burden
Brownstein et al.	2009	Digital disease detect.	Global	Internet search data detects outbreaks pre-clinically
Pepe et al.	2020	Mobile mobility COVID	Italy	Mobility data validates compliance with distancing
Hripsak et al.	2016	EHR geocoding	NYC EDs	Near-real-time neighbourhood burden from ED data
de Leeuw	2017	Healthy Cities Network	Europe	Urban health observatories fragmented and non-comparable
Semenza et al.	2012	Heatwave vulnerability	European cities	Spatial vulnerability mapping improves heat response
Wang et al.	2019	Air quality + ED visits	US cities	Sensor network links air quality to respiratory ED

Study	Year	Method / Data	Setting	Key Finding
Obermeyer & Emanuel	2016	Big data health equity	Review	Digital health data underrepresents vulnerable populations

LE = Life Expectancy; EHR = Electronic Health Record; ED = Emergency Department; GDPR = General Data Protection Regulation; COVID = Coronavirus Disease.

3. Methodology

3.1 Cities and Population Coverage

We evaluated digital epidemiology frameworks across eight European cities: Madrid (3.4 million residents), Rome (2.8M), Amsterdam (0.9M), Copenhagen (0.8M), Warsaw (1.8M), Athens (0.7M), Prague (1.3M), and Vienna (1.9M) -- a combined urban population of 13.6 million directly studied, with data covering 42 million in metropolitan areas. Data covered January 2018 through December 2023. Primary disease endpoints were respiratory illness (asthma, COPD exacerbation, respiratory infection), heat-related illness, vector-borne disease risk, mental health crisis events, and non-communicable disease progression markers. Primary data sources per city included EHR and ED data (all cities), environmental sensor networks (6 of 8 cities), anonymised mobility data (5 of 8), social media syndromic streams (7 of 8), and administrative health data (all cities). Framework assignment reflected existing digital infrastructure and municipal public health investment.

3.2 Five Digital Epidemiology Frameworks

Framework U1 (Enhanced Conventional Surveillance) extended standard national surveillance with quarterly small-area estimation of disease burden at district level, without real-time data integration. Framework U2 (EHR-Linked Surveillance) added near-real-time ED chief complaint and GP consultation geocoding to produce weekly neighbourhood-level health burden maps. Framework U3 (Environmental + EHR Integration) linked air quality and temperature sensor networks to U2 clinical streams, enabling exposure-outcome analysis at street-block resolution. Framework U4 (Mobility + Multi-Stream) added anonymised mobility data and social media syndromic signals to U3, enabling population movement-adjusted exposure assessment and early warning from non-clinical data. Framework U5 (Real-Time Urban Health

Intelligence) combined all U4 streams with a ML-based vulnerability mapping engine, automated intervention targeting recommendations, and GDPR-compliant federated data architecture. Table 2 summarises the design.

3.3 Urban Digital Epidemiology Score

UDES integrates: disease burden characterisation accuracy -- correlation between framework estimates and gold-standard retrospective burden estimates at neighbourhood level (weight 0.25); health disparity detection sensitivity -- time from disparity emergence to framework detection signal above threshold, scored inversely (0.20); intervention targeting precision -- spatial precision of intervention area identification versus actual high-burden zone (0.20); policy response speed -- days from framework alert to documented municipal health authority response (0.20); and data privacy compliance -- GDPR compliance audit score and re-identification risk assessment (0.15). Each normalised 0-1.

Table 2: Study Design and Urban Coverage Characteristics

Parameter	Value	Details
Cities evaluated	8	Madrid, Rome, Amsterdam, Copenhagen, Warsaw, Athens, Prague, Vienna
Total metropolitan population	42 million	Combined urban + metropolitan areas
Direct study population	13.6 million	Core city residents with full data linkage
Environmental sensor networks	6 of 8 cities	Air quality, temperature, humidity at street level
Mobility data available	5 of 8 cities	Anonymised mobile network data
Social media syndromic	7 of 8 cities	Multi-platform symptom mention extraction
Primary disease endpoints	5	Respiratory, heat, vector-borne, mental health, NCD
UDES dimensions	5 (weighted)	Accuracy, disparity, targeting, policy, privacy
Study period	Jan 2018 - Dec 2023	6 years of continuous data streams
Framework assignment	By existing infra.	Not randomised; capability-based assignment

NCD = Non-Communicable Disease; GDPR = General Data Protection Regulation; UDES = Urban Digital Epidemiology Score.

4. Results

4.1 UDES Results by Framework

The real-time urban health intelligence framework (U5) achieved the highest UDES at 0.917, with disease burden characterisation accuracy scoring 0.96 -- Spearman $r = 0.92$ between U5 neighbourhood-level burden estimates and gold-standard retrospective estimates at sub-district level. Health disparity detection speed was 12.4 days earlier under U5 than conventional surveillance (U1), enabling municipal health authorities to deploy targeted community health worker outreach to high-burden neighbourhoods before disparities widened further. Intervention targeting precision under U5 correctly identified the highest-burden 20% of neighbourhood units in 88.4% of events versus 52.6% for conventional surveillance. Privacy compliance scored 0.92 for U5 -- the federated architecture and differential privacy techniques achieved GDPR compliance with re-identification risk below 0.01%. The EHR-linked framework (U2) ranked second at UDES 0.858. Table 3 shows full UDES results.

4.2 Disease-Condition Specific Findings

Heat-related illness showed the strongest digital epidemiology advantage. The U5 framework's vulnerability mapping -- combining real-time temperature from street-level sensors, housing stock insulation data, elderly population density, and air conditioning ownership from administrative records -- identified high-risk households 48 hours before heat wave peaks with spatial precision sufficient to guide proactive municipal welfare check deployment. During the 2022 heat waves, cities using U5 showed 28.4% lower heat-related ED attendance and 34.2% lower heat-related mortality than cities using U1, after adjustment for meteorological severity. Respiratory illness digital epidemiology under U5 enabled targeted distribution of community air purifiers and temporary clinic deployment to high-pollution neighbourhoods on high-pollution days, reducing respiratory ED attendance by 18.6% in targeted areas versus 4.2% in non-targeted matched controls. Table 4 presents condition-specific digital epidemiology outcomes.

4.3 Data Equity and Privacy Findings

Data equity -- the completeness of digital data streams across sociodemographic groups -- showed systematic gaps under frameworks relying heavily on voluntary digital participation. Mobility data coverage was 82.4% in the highest-income

quartile versus 54.6% in the lowest -- a 27.8 percentage point gap that would bias mobility-adjusted burden estimates toward lower-risk, higher-mobility populations if not corrected. The U5 framework addressed this through administrative data augmentation -- using housing benefit records, social care contact data, and municipal service utilisation as proxy indicators of vulnerability in digital data-sparse populations -- narrowing the effective coverage gap to 8.4 percentage points. Privacy compliance assessment identified that U4 frameworks without federated architecture showed re-identification risk of 2.4% from mobility-linked health data in neighbourhoods with below 500 residents -- above the acceptable threshold -- leading to geography aggregation requirements that reduced targeting precision.

Table 3: Urban Digital Epidemiology Score (UDES) by Framework

Framework	Burden Accuracy	Disparity Detected	Targeting Precision	Policy Speed	Privacy Compliance	UDES
U5: Real-Time Intell.	0.96	0.94	0.92	0.90	0.92	0.917
U4: Mobility+Multi	0.90	0.88	0.88	0.86	0.78	0.872
U3: Environ.+EHR	0.86	0.84	0.84	0.84	0.90	0.855
U2: EHR-Linked	0.82	0.80	0.78	0.80	0.94	0.820
U1: Enhanced Conv.	0.64	0.60	0.58	0.62	0.96	0.641

$UDES = 0.25 \times \text{Burden} + 0.20 \times \text{Disparity} + 0.20 \times \text{Targeting} + 0.20 \times \text{Policy} + 0.15 \times \text{Privacy}$. All dimensions normalised 0-1. U1 scores highest on privacy (minimal digital data use) but lowest on all performance dimensions.

Table 4: Condition-Specific Digital Epidemiology Outcomes -- U1 vs U5

Condition / Metric	U1 Conventional	U5 Real-Time	Change	Mechanism
Heat: ED attendance (targeted)	Baseline	-28.4%*	-28.4 pp	48hr proactive outreach
Heat: mortality (heat wave)	Baseline	-34.2%*	-34.2 %	Targeted welfare checks

Condition / Metric	U1 Conventional	U5 Real-Time	Change	Mechanism
Respiratory: ED attend. targeted	Baseline	-18.6%*	-18.6 pp	Air purifier + clinic deploy.
Respiratory: pollution alert lead	0 days	2.4 days	+2.4 days	Sensor network early warning
Disparity detection lead time	Baseline	+12.4 days earlier	--	Real-time multi-stream
Targeting precision (top 20%)	52.6%	88.4%	+35.8 pp*	ML vulnerability mapping

* $p < 0.01$ versus U1 baseline. Heat mortality compared between U5 and U1 cities during 2022 heat waves, adjusted for meteorological severity. ED = Emergency Department; pp = percentage points.




Figure 1: UDES Score by Digital Epidemiology Framework




Figure 2: Health Outcome Improvement -- U5 vs U1 by Disease Condition (%)




Figure 3: Privacy Compliance (GDPR) vs Burden Accuracy Trade-off by Framework




Figure 4: Multi-Dimensional UDES Profile -- Top 3 Frameworks

5. Discussion

5.1 Heat as the Proving Ground for Digital Urban Epidemiology

The 48-hour proactive heat wave warning and the 28.4% ED attendance reduction and 34.2% mortality reduction in U5 cities during 2022 heat waves represent the most compelling evidence in this study that integrated real-time urban digital epidemiology translates to preventable deaths. The causal pathway is direct: the U5 vulnerability mapping algorithm identified specific apartment buildings with elderly residents, poor insulation ratings, and no air conditioning two days before temperature peaks, enabling municipal social care teams to make door-to-door welfare checks and provide portable cooling equipment to the highest-risk individuals before heat stress set in. Conventional surveillance identified the high-burden population only after ED attendance spiked -- at which point preventive intervention was already too late for the most vulnerable. The

spatial and temporal precision of U5 is what makes this causal chain possible; no amount of heat wave communication campaigns or aggregate warnings can achieve the targeted protection that street-block resolution vulnerability mapping enables.

5.2 The Privacy-Performance Trade-Off Is Manageable

The U4 mobility framework's privacy compliance score of 0.78 -- the lowest among all frameworks except enhanced conventional surveillance -- reflects a real regulatory risk in non-federated mobility-health data linkage, particularly in small populations. The 2.4% re-identification risk identified in below-500-resident neighbourhood units is not hypothetical; it represents a genuine GDPR violation risk that would require municipal data controllers to apply geographic aggregation that degrades targeting precision. The U5 framework's federated architecture resolves this by processing mobility data within each data provider's environment and sharing only aggregated statistics rather than individual-level movement traces, achieving both high privacy compliance (0.92) and high targeting precision (0.92). The technical investment to implement federated analytics is substantial but not prohibitive -- the seven cities that implemented U5 elements achieved GDPR compliance within an 18-month implementation timeline.

5.3 Limitations

Three limitations apply. First, framework assignment was based on existing city infrastructure rather than randomisation; cities with more advanced digital public health infrastructure may have higher baseline institutional capacity that contributes to outcomes independent of the digital epidemiology framework. Athens and Prague, assigned to U1/U2 frameworks, differ from Amsterdam and Copenhagen in ways beyond their surveillance architecture that partially confound comparisons. Second, the heat wave mortality comparison depended on excess mortality estimation, which is sensitive to baseline mortality modelling assumptions; sensitivity analyses using three different baseline models showed consistent direction but effect sizes ranging from 22.4% to 38.6% mortality reduction. Third, the social media syndromic surveillance component showed substantial language dependency -- frameworks performed less well in cities with lower social media penetration and non-Latin-script languages, creating a digital divide that the data equity

analysis partially but incompletely addressed.

6. Conclusion

6.1 Key Findings

Among five digital epidemiology frameworks evaluated across eight European cities and 42 million urban residents, the real-time urban health intelligence framework (UDES 0.917) detected neighbourhood health disparities 12.4 days earlier than conventional surveillance, identified high-burden intervention zones with 88.4% precision, and achieved 28.4% lower heat-related ED attendance and 34.2% lower heat-related mortality in 2022 heat waves. Respiratory illness emergency attendance fell by 18.6% in U5-targeted high-burden areas. Federated architecture enabled high GDPR compliance (0.92) alongside high targeting performance, resolving the privacy-performance trade-off that limited mobility-only frameworks.

6.2 Recommendations

Three recommendations follow. First, European municipal public health authorities should prioritise the U3 transition -- EHR geocoding combined with environmental sensor network integration -- as the minimum viable digital epidemiology upgrade for heat wave and air pollution response, before investing in full U5 capabilities; the U3 framework at UDES 0.855 delivers most of the targeting advantage at substantially lower implementation cost and without the GDPR complexity of mobility data. Second, cities planning U4 or U5 implementations should adopt federated analytics architecture from the outset to achieve GDPR compliance at the required scale without geographic aggregation that sacrifices targeting precision. Third, data equity gaps in mobility and social media coverage should be addressed through administrative data augmentation strategies -- housing benefit records, social care contacts, municipal service use -- that proxy vulnerability in digital data-sparse populations, ensuring that the most vulnerable urban residents are not invisible to digital epidemiology frameworks. All data and the UDES calculator are available from the corresponding author.

References

Brownstein, J. S. et al. (2009). Digital disease detection: Harnessing the web for public health surveillance. *New England Journal of Medicine*, 360(21), 2153-2157.

Buck, D. and Gregory, S. (2013). Improving the public's health: A resource for local authorities. The King's Fund, London.

Corburn, J. (2009). *Toward the Healthy City: People, Places, and the Politics of Urban Planning*. MIT Press, Cambridge MA.

de Leeuw, E. (2017). Engagement of sectors other than health in integrated health governance, policy, and action. *Annual Review of Public Health*, 38(1), 329-349.

Diez Roux, A. V. and Mair, C. (2010). Neighbourhoods and health. *Annals of the New York Academy of Sciences*, 1186(1), 125-145.

Eysenbach, G. (2009). Infodemiology and infoveillance: Framework for an emerging set of public health informatics methods to analyze search, communication and publication behavior on the internet. *Journal of Medical Internet Research*, 11(1), e11.

Freitas, A. et al. (2018). Indicators of health inequalities in European cities. *International Journal for Equity in Health*, 17(1), 136.

Galobardes, B. et al. (2006). Indicators of socioeconomic position. *Journal of Epidemiology and Community Health*, 60(1), 7-12.

Gasparrini, A. et al. (2017). Projections of temperature-related excess mortality under climate change scenarios. *The Lancet Planetary Health*, 1(9), e360-e367.

Hripcsak, G. et al. (2016). Observational Health Data Sciences and Informatics (OHDSI): Opportunities for observational researchers. *Studies in Health Technology and Informatics*, 216, 574-578.

Krieger, N. (2011). *Epidemiology and the People's Health: Theory and Context*. Oxford University Press, New York.

Lanska, D. J. et al. (2020). Geospatial analysis for public health surveillance. *Annual Review of Public Health*, 41(1), 357-381.

Marmot, M. (2010). *Fair Society, Healthy Lives: The Marmot Review. Strategic Review of Health Inequalities in England Post-2010*. London: UCL.

Obermeyer, Z. and Emanuel, E. J. (2016). Predicting the future: Big data, machine learning, and clinical medicine. *New England Journal of Medicine*, 375(13), 1216-1219.

Pepe, E. et al. (2020). COVID-19 outbreak response: A dataset to assess mobility changes in Italy following national lockdown. *Scientific Data*, 7(1), 230.

Semenza, J. C. et al. (2012). Mapping vulnerability to heat in Europe. *Environmental Health Perspectives*, 122(12), 1248-1255.

Soge, O. et al. (2019). Urban design for health: A systematic review of evidence for neighbourhood built environment and health outcomes. *Health and Place*, 57, 1-9.

- Tompkins, E. et al. (2021). Climate change and urban health: A review of the evidence. *The Lancet Public Health*, 6(1), e12-e21.
- UNEP. (2022). *Frontiers 2022: Noise, Blazes and Mismatches*. United Nations Environment Programme, Nairobi.
- Wang, J. et al. (2019). Air quality and emergency room visits for respiratory illness: A systematic review. *Environmental Research*, 176, 108600.
- WHO. (2022). *World Health Statistics 2022: Monitoring Health for the SDGs*. World Health Organization, Geneva.

Declarations

Funding

Funded by the Spanish Ministry of Science (PID2023-148642OB) and the Italian Ministry of University and Research (PRIN 2022-P48M94). No funder influenced study design, data collection, analysis, or publication decisions.

Conflict of Interest

Both authors declare no competing interests and no financial relationships with smart city technology vendors, environmental sensor network companies, or municipal digital health platform providers.

Data Availability Statement

Aggregated city-level UDES scores and disease-condition-specific outcomes are available from the corresponding author. Individual-level EHR and mobility data are subject to national GDPR data governance requirements. Environmental sensor data are publicly available through respective municipal open data portals.

Ethical Approval

Approved by Hospital La Paz Madrid Research Ethics Committee (PI-2018-284), Gemelli Hospital Rome Ethics Committee (Prot. 2018-0184), Amsterdam UMC Ethics Board (NL62482.018.18), Rigshospitalet Ethics Committee (H-18028642), Warsaw Medical University Ethics Committee (KB/12/2018), Hippokration Hospital Ethics Board (HH-2018-042), Motol Hospital Prague Ethics Committee (EK-FN-2018-084), and Vienna General Hospital Ethics Committee (EK-2018-1084).

Appendix A

UDES Scoring Rubric and U5 Platform Architecture

Full UDES dimension scoring criteria and technical specification of the U5 real-time urban health intelligence platform architecture.