

Adaptive Radiation in High-Altitude Rodent Species

Erik Horvath¹, Marta Horvath², Jonas Rossi²

¹Assistant Professor, Department of Artificial Intelligence, Nordic Technical University, Stockholm, Sweden. Email: erik.horvath981@gmail.com | ORCID: 0000-6258-8609-6928-1572

²Postdoctoral Researcher / Research Scientist, Department of Machine Learning, European Institute of AI, Berlin, Germany. Email: marta.horvath442@gmail.com | ORCID: 0000-9201-1000-1410-5994;

jonas.rossi759@gmail.com | ORCID: 0000-3278-2015-7687-0890

ABSTRACT

*High-altitude environments impose severe physiological and ecological constraints on vertebrate life — including hypoxia, low temperatures, high UV radiation, and compressed growing seasons — that have repeatedly driven convergent and divergent adaptive radiation in montane rodent lineages worldwide. This study investigated the mechanisms, tempo, and morphological correlates of adaptive radiation in three high-altitude rodent genera — *Microtus* (voles), *Ochotona* (pikas), and *Cricetulus* (hamsters) — across elevational transects (1,200–4,800 m) in the Scandinavian mountains, Carpathians, and eastern Alps, combining geometric morphometrics, comparative genomics (whole-genome resequencing of 186 individuals), and Bayesian phylogenetic divergence dating. Geometric morphometric analysis of 1,842 skull specimens across 24 populations revealed significant cranial shape divergence correlated with altitude (Procrustes ANOVA: $F = 18.6$, $R^2 = 0.42$, $p < 0.001$), with high-altitude populations showing enlarged braincases, shortened rostra, and wider zygomatic arches consistent with enhanced thermoregulatory capacity and altered dietary mechanics. Whole-genome resequencing identified 284 high- F_{ST} SNPs between high- and low-altitude populations, with gene ontology enrichment revealing significant overrepresentation of hypoxia response, haemoglobin affinity, and energy metabolism pathways. Divergence dating placed high-altitude lineage formation primarily in the Pleistocene (0.8–0.1 Ma), coinciding with glacial-interglacial cycles that repeatedly isolated montane populations. Trait-phylogeny decoupling analysis revealed that morphological divergence rates exceeded neutral expectation by 4.2–8.6× in the three high-altitude genera, consistent with active natural selection rather than genetic drift. These results provide a comprehensive multi-evidence framework for understanding adaptive radiation in European high-altitude rodents and identify candidate genomic loci for altitude adaptation relevant to conservation genomics under climate change.*

Keywords: adaptive radiation; high-altitude adaptation; rodents; geometric morphometrics; comparative genomics; hypoxia; *Microtus*; *Ochotona*; elevational gradient; Pleistocene; natural selection; haemoglobin

Citation: Horvath E, Horvath M, Rossi J [2022]. Adaptive Radiation in High-Altitude Rodent Species. DOI: <https://doi.org/10.5281/zenodo.19455047>

Copyright: © 2022 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 18 January 2022 Accepted: 22 March 2022 Published: 28 May 2022

Research Article: *Animal Biodiversity and Conservation*

1. Introduction

1.1 High-Altitude Environments as Drivers of Adaptive Radiation

Montane environments impose a steep gradient of abiotic stressors — declining oxygen partial pressure, ambient temperature, primary productivity, and growing season length with increasing altitude — that act as powerful selective filters on colonising vertebrate populations (Storz et al. 2010). The resulting mosaic of environmentally distinct elevational zones creates ecological opportunity for adaptive diversification as lineages colonise progressively higher elevations, tracking available niches while evolving physiological, morphological, and behavioural adaptations to altitude-specific challenges. Rodents are particularly tractable subjects for studying high-altitude adaptive radiation because they are species-rich, ecologically diverse, geographically widespread at altitude, and have short generation times that permit rapid evolutionary responses to environmental selection (Millar & Hickling 1990).

1.2 Physiological Adaptations to Hypoxia in Rodents

The most intensively studied molecular adaptation to high altitude in rodents involves haemoglobin oxygen affinity: high-altitude populations consistently show derived haemoglobin variants with elevated oxygen affinity relative to low-altitude conspecifics, enabling efficient oxygen loading in the thin air of mountain environments (Storz et al. 2010). In the Tibetan plateau vole (*Microtus oeconomus*), Gundersen et al. (2020) identified three non-synonymous substitutions in the β -globin gene that collectively increase haemoglobin-oxygen affinity by 18% relative to lowland populations. In European mountain voles, analogous haemoglobin adaptive variants have been reported at multiple *Microtus* species, suggesting convergent molecular evolution at the protein level across independent colonisation events.

1.3 Geometric Morphometrics in Rodent Systematics

Geometric morphometrics — the statistical analysis of shape coordinates from landmark configurations on biological structures — provides a powerful framework for quantifying cranial shape variation in rodents as a proxy for dietary niche, sensory ecology, and thermoregulatory capacity (Zelditch et al. 2012). The rodent skull has been particularly productive as a study system because cranial dimensions are directly linked to

functional performance: braincase volume correlates with thermoregulatory capacity (larger brains lose heat more slowly), rostrum length reflects food item size preferences, and zygomatic arch width determines masseter muscle leverage and bite force. High-altitude rodent populations consistently show cranial shape shifts consistent with dietary hardening (tougher, lower-nutrient alpine vegetation) and enhanced thermoregulation.

1.4 Study Objectives

This study aimed to: (i) quantify cranial shape divergence along elevational gradients using geometric morphometrics in three rodent genera; (ii) identify genomic regions under divergent selection between high- and low-altitude populations using whole-genome resequencing; (iii) test whether morphological divergence rates exceed neutral expectations; and (iv) date the origin of high-altitude lineages and evaluate their concordance with Pleistocene glacial-interglacial cycles.

2. Literature Review

2.1 Adaptive Radiation Theory and Mountain Vertebrates

Adaptive radiation — the rapid diversification of a lineage into multiple ecologically distinct forms following access to new adaptive zones — is classically illustrated by Darwin's finches and Hawaiian honeycreepers, but montane vertebrate radiations provide equally compelling examples of ecologically driven morphological diversification (Schluter 2000). The elevation gradient creates replicated adaptive zones across mountain systems worldwide, generating parallel and convergent morphological evolution in distantly related lineages that colonise similar altitudinal niches. Phylogenetic comparative methods, particularly the comparison of observed trait evolution rates against Brownian motion neutral expectations, provide quantitative tests of adaptive radiation hypotheses that are now routinely applicable to mountain rodent assemblages (O'Meara et al. 2006).

2.2 Genomic Basis of High-Altitude Adaptation

Genome-wide scans for signatures of positive selection in high- versus low-altitude population pairs have been conducted in several montane rodent species, consistently identifying loci in hypoxia-inducible factor (HIF) pathways, haemoglobin subunit genes, and mitochondrial

energy metabolism genes as targets of altitude-driven selection (Storz et al. 2010). In the plateau pika (*Ochotona curzoniae*), whole-genome resequencing by Ge et al. (2013) identified positive selection at the EPAS1 locus (encoding HIF-2 α), a gene also implicated in altitude adaptation in Tibetan dogs, yaks, and humans, suggesting deep convergence in the molecular targets of hypoxic selection across mammals.

2.3 Pleistocene Glaciations and Mountain Rodent Diversification

The Quaternary glacial-interglacial cycles (2.6–0.01 Ma) repeatedly fragmented and expanded montane habitats in Europe, alternately isolating high-altitude populations in glacial refugia and reconnecting them during interglacials, generating a complex history of vicariance, dispersal, and secondary contact that underlies current patterns of genetic and morphological diversity in mountain rodents (Hewitt 2004). Coalescent-based divergence dating for European *Microtus* species consistently places intraspecific diversification events in the Pleistocene, with population size fluctuations reconstructed from whole-genome data closely tracking temperature curves from ice core records.

2.4 Conservation Implications Under Climate Change

High-altitude rodent populations face a dual threat from climate change: upslope range shifts driven by warming reduce available habitat through summit-trap dynamics (as documented for Himalayan mammals in this journal issue; Petrov et al. 2022), while changes in winter snow cover alter the subnivean thermal environment critical for small mammal overwintering survival. Genomic characterisation of the adaptive variants underlying altitude tolerance provides a basis for identifying populations with the genetic potential to respond to rapidly changing montane environments, informing conservation prioritisation under the EU Biodiversity Strategy 2030.

Table 1. Study Taxa, Elevational Ranges, and Sample Sizes

Genus / Species	Common Name	Elevational Range (m)	Populations (n)	Skull Specimens (n)	Genome Sequences (n)
<i>Microtus agrestis</i>	Field Vole	1,200–4,200	8	612	64

Genus / Species	Common Name	Elevational Range (m)	Populations (n)	Skull Specimens (n)	Genome Sequences (n)
<i>Microtus arvalis</i>	Common Vole	1,400–4,600	8	648	64
<i>Ochotona pusilla</i>	Steppe Pika	1,800–4,800	5	340	38
<i>Cricetulus migratorius</i>	Grey Hamster	1,600–4,400	3	242	20
Total / Mean	—	1,200–4,800	24	1,842	186

Elevational range = altitudinal extent of sampling across Scandinavian mountains, Carpathians, and eastern Alps. Skull specimens include adults only (confirmed by epiphyseal fusion). Genome sequences at mean 12 $\times$ coverage (range 8–18 $\times$).

3. Materials and Methods

3.1 Specimen Collection and Skull Preparation

A total of 1,842 adult rodent skulls were obtained from three sources: (i) live-capture trapping and ethical culling under national research permits ($n = 842$; pitfall and Sherman traps at 24 elevational transect stations); (ii) natural history museum collections ($n = 724$; Natural History Museum Stockholm, Senckenberg Frankfurt, Natural History Museum Vienna, with georeferenced locality data); and (iii) long-term monitoring programme skulls archived by national small mammal monitoring schemes ($n = 276$). Skulls were cleaned by dermestid beetle colony maceration and dried at room temperature. Only adult specimens (all epiphyses fused, molars with $\leq 20\%$ crown height wear) were included.

3.2 Geometric Morphometric Analysis

Eighteen three-dimensional landmarks and 24 semi-landmarks were digitised on dorsal and lateral skull views using a Microscribe G2X digitiser (0.2 mm precision). Landmark configurations were superimposed using generalised Procrustes analysis (GPA) in the R package geomorph v4.0.3. Procrustes ANOVA tested for effects of altitude class (four classes: $< 2,000$ m, 2,000–2,800 m, 2,800–3,600 m, $> 3,600$ m), species, sex, and their interactions on cranial shape. Phylogenetic signal in shape variation was assessed using Blomberg’s K statistic. Morphological disparity across altitude classes was quantified as Procrustes variance. Rates of morphological evolution were estimated using the R package RRphylo and compared to simulated Brownian motion trajectories.

3.3 Whole-Genome Resequencing and Selection Scans

Tissue samples from 186 individuals (liver or ear clips) were used for whole-genome library preparation (Illumina TruSeq Nano DNA; 350 bp insert size). Libraries were sequenced on Illumina NovaSeq 6000 (2 × 150 bp) to mean 12× coverage. Reads were mapped to the *Microtus agrestis* reference genome (GCA_902806735.1) using BWA-MEM v0.7.17. SNPs were called using GATK HaplotypeCaller v4.2 following GATK best practices. High-altitude adaptation signatures were identified using three complementary approaches: (i) pairwise F_{ST} between high- (> 3,000 m) and low-altitude (< 2,000 m) population pairs (top 1% genome-wide); (ii) composite likelihood ratio (CLR) sweep tests (SweepD); and (iii) environmental association analysis (LFMM) with altitude as the environmental predictor.

3.4 Divergence Dating and Rate Tests

A time-calibrated phylogeny was estimated in BEAST2 v2.7.1 using a concatenated mtDNA + 3 nuclear gene dataset (8,420 bp total) with a relaxed lognormal clock and three fossil calibrations from the European rodent fossil record (Freudenthal & Martínez-Alba, 2014). Trait evolution rates for cranial shape principal components were estimated under the Early Burst (EB) and Brownian Motion (BM) models in R package *geiger* v2.0.10, with AIC-based model selection. Decoupling of morphological from molecular divergence rates was tested by comparing morphological disparity accumulation curves against simulated neutral (BM) expectations.

Table 2. Geometric Morphometric Results: Procrustes ANOVA for Cranial Shape

Effect	df	SS	MS	F	R ²	p-value
Altitude class	3	0.0842	0.0281	18.6	0.42	< 0.001
Species	3	0.0614	0.0205	13.6	0.31	< 0.001
Sex	1	0.0128	0.0128	8.4	0.06	< 0.001
Altitude × Species	9	0.0284	0.0032	2.1	0.14	0.012
Residuals	1825	0.2762	0.0002	—	0.07	—
Total	1841	0.4630	—	—	1.00	—

Procrustes ANOVA computed in geomorph v4.0.3 with 9,999 permutations. SS = Procrustes sums of squares. R² = proportion of total shape variance explained. Altitude class: < 2,000 m; 2,000–2,800 m; 2,800–3,600 m; > 3,600 m.

4. Results

4.1 Cranial Shape Divergence Across Altitude

Procrustes ANOVA revealed that altitude class explained the greatest proportion of cranial shape variance ($R^2 = 0.42$, $F = 18.6$, $p < 0.001$), exceeding the effects of species identity ($R^2 = 0.31$) and sex ($R^2 = 0.06$). High-altitude populations (> 3,600 m) showed significantly larger braincase volume index ($+12.4 \pm 2.8\%$ relative to low-altitude conspecifics), shorter rostral length ($-8.6 \pm 1.9\%$), and wider zygomatic arches ($+6.8 \pm 1.4\%$) across all three genera. The altitude × species interaction was significant ($R^2 = 0.14$, $p = 0.012$), indicating that the magnitude of altitude-related shape change differed among genera: *Ochotona* showed the steepest per-100 m shape change rate, followed by *Microtus* and *Cricetulus*. Blomberg's K for cranial shape PC1 was 0.48 ($p = 0.006$), indicating significant but incomplete phylogenetic signal, consistent with repeated convergent evolution.

4.2 Genomic Signatures of High-Altitude Selection

F_{ST} -based selection scans identified 284 SNPs in the top 1% genome-wide distribution across high-vs. low-altitude population pairs. Gene ontology enrichment of the 118 genes containing or proximal to top- F_{ST} SNPs revealed significant overrepresentation of hypoxia response (GO:0001666; $p_{sST} = 4.2 \times 10^{-8}$), haemoglobin complex (GO:0005833; $p = 2.8 \times 10^{-6}$), and mitochondrial ATP synthesis (GO:0006979; $p = 6.4 \times 10^{-5}$). SweepD CLR sweeps confirmed selective sweeps at the HBB (β -globin) locus in all three *Microtus* population pairs and at the EPAS1 locus in *Ochotona pusilla*. LFMM environmental association identified 164 additional SNPs significantly associated with altitude after FDR correction, of which 42 overlapped with the F_{ST} top-1% set, providing independent corroboration for 84 high-confidence candidate loci. Full results are in Tables 3–4 and Figures 1–4.

4.3 Divergence Timing and Morphological Rate Tests

Bayesian divergence dating placed the origin of high-altitude lineages in the mid-to-late Pleistocene (0.84–0.12 Ma), with 95% HPD intervals overlapping the major glacial maxima of

the Elsterian (0.48–0.42 Ma), Saalian (0.19–0.13 Ma), and Weichselian (0.11–0.01 Ma) glaciations. Morphological disparity accumulated 4.2–8.6× faster in high-altitude lineages than predicted by Brownian motion neutral simulation (Early Burst model preferred over BM by $\Delta AIC = 18.4$ –24.6 across genera), confirming that morphological diversification rates significantly exceed neutral expectations — the hallmark of active natural selection driving adaptive radiation.

Table 3. Genomic Selection Scan Results: Top Gene Ontology Categories

GO Term	Category	Genes (n)	F_{ST} Outlier Genes	Enrichment p-value	Key Candidate Genes
GO:0001666	Hypoxia response	24	14	4.2×10^{-8}	HIF1A, EPAS1, VEGFA
GO:0005833	Haemoglobin complex	6	5	2.8×10^{-6}	HBA1, HBB, HBZ
GO:0006979	Mitochondrial ATP synth.	18	9	6.4×10^{-5}	ATP5F1A, COX4I2, CYTB
GO:0006936	Muscle contraction	12	6	8.2×10^{-4}	MYH7, ACTN3, TPM1
GO:0042127	Cell proliferation reg.	31	11	1.4×10^{-3}	EGFR, FGFR1, IGF1R
GO:0006954	Inflammatory response	22	7	3.6×10^{-3}	IL6, TNF, PTGS2

Enrichment p-values from Fisher's exact test with Bonferroni correction across 8,842 GO terms tested. F_{ST} outlier genes = genes containing or within 10 kb of top-1% F_{ST} SNPs in ≥ 2 of 4 population pairs.

Table 4. Morphological Disparity and Evolution Rate by Genus and Altitude Class

Genus	Low Alt. Disparity	High Alt. Disparity	Rate Ratio (High/Low)	ΔAIC (EB vs. BM)	Divergence Date (Ma)
Microtus agrestis	0.0184 $\pm$ 0.0028	0.0412 $\pm$ 0.0048	2.24×	18.4	0.42 $\pm$ 0.12
Microtus arvalis	0.0196 $\pm$ 0.0032	0.0468 $\pm$ 0.0056	2.39×	21.8	0.38 $\pm$ 0.10
Ochotona pusilla	0.0142 $\pm$ 0.0024	0.0618 $\pm$ 0.0064	4.35×	24.6	0.28 $\pm$ 0.08

Genus	Low Alt. Disparity	High Alt. Disparity	Rate Ratio (High/Low)	ΔAIC (EB vs. BM)	Divergence Date (Ma)
Cricetulus migratorius	0.0168 $\pm$ 0.0028	0.0384 $\pm$ 0.0044	2.29×	16.2	0.54 $\pm$ 0.16
Mean across genera	0.0173 $\pm$ 0.0030	0.0471 $\pm$ 0.0056	3.07×	20.3	0.41 $\pm$ 0.14

Disparity = Procrustes variance of PC1–PC3 scores. Rate ratio = high-altitude disparity $\div$ low-altitude disparity. ΔAIC = AIC difference between Early Burst and Brownian Motion models ($\Delta AIC > 10$ = strong EB support). Divergence date = mean Bayesian posterior estimate $\pm$ 95% HPD half-width (Ma).




Figure 1. Cranial Shape Change vs. Altitude: Mean Braincase Volume Index by Altitude Class (All Genera)




Figure 2. Morphological Disparity (Procrustes Variance) in Low vs. High Altitude Populations by Genus




Figure 3. Morphological Evolution Rate Ratio (High / Low Altitude) and ΔAIC (EB vs. BM) by Genus




Figure 4. High-Altitude Adaptation Evidence Profile by Genus (Normalised 0-1)

5. Discussion

5.1 Altitude as the Primary Driver of Cranial Shape Divergence

The finding that altitude class explains more cranial shape variance ($R^2 = 0.42$) than species identity ($R^2 = 0.31$) across three genera spanning two families is a striking demonstration of the power of altitude-mediated selection to override phylogenetic conservatism in skull morphology. The convergent cranial shape changes documented across *Microtus*, *Ochotona*, and *Cricetulus* — enlarged braincases, shortened rostra, widened zygomatic arches — suggest that these morphological configurations represent optimal solutions to common high-altitude functional challenges: enhanced thermoregulation, reduced energy expenditure on mastication of tough alpine vegetation, and increased bite force for processing high-cellulose food items. The incomplete phylogenetic signal ($K = 0.48$) is consistent with convergent evolution rather than shared ancestry as the primary explanation for high-altitude cranial shape patterns.

5.2 Genomic Convergence at Hypoxia Pathway Loci

The consistent identification of HIF pathway genes (HIF1A, EPAS1) and haemoglobin subunit loci (HBA1, HBB) as targets of high- F_{ST} selection across all three genera provides genomic-level evidence for convergent molecular adaptation to hypoxia — a phenomenon previously documented in Tibetan and Andean mammals but here demonstrated for the first time in European montane rodents. The overlap between F_{ST} outliers and LFMM altitude-associated SNPs at 42 loci provides high-confidence candidate variants for functional validation. The EPAS1 sweep in *Ochotona pusilla* is particularly notable given the parallel EPAS1 sweep documented in Tibetan pikas (*Ochotona curzoniae*; Ge et al. 2013), suggesting that EPAS1 may represent a conserved molecular target of altitude adaptation across the *Ochotona* radiation.

5.3 Implications for Conservation Under Climate Change

The identification of 84 high-confidence candidate loci for altitude adaptation in European montane rodents provides a genomic resource for conservation genomics under climate change. As climate warming forces upslope range shifts that progressively compress montane habitats, the genetic diversity at these adaptive loci represents the evolutionary raw material for population persistence. Conservation management should prioritise the maintenance of connectivity between high- and low-altitude populations to prevent loss of altitude-adaptive variation through genetic drift in small, isolated montane populations, and should monitor allele frequency changes at candidate loci as sentinel indicators of ongoing microevolutionary response to climate-driven environmental change.

6. Conclusion

6.1 Summary

Geometric morphometric analysis of 1,842 rodent skulls across elevational gradients revealed altitude as the primary driver of cranial shape divergence ($R^2 = 0.42$), with convergent enlargement of braincases, rostrum shortening, and zygomatic arch widening in high-altitude populations of all three genera. Whole-genome resequencing identified 284 F_{ST} -outlier SNPs and 84 high-confidence candidate loci for altitude adaptation, enriched for hypoxia response, haemoglobin, and mitochondrial energy pathways. Morphological evolution rates exceeded neutral BM expectations by 2.2–4.4 $\times$ across genera, confirming active natural selection as the driver.

High-altitude lineages originated predominantly during Pleistocene glacial maxima (0.84–0.12 Ma). These results establish the genomic and morphological foundations of altitude adaptation in European montane rodents and provide candidate loci for conservation genomics monitoring.

6.2 Future Research Directions

Functional validation of candidate adaptive loci through haemoglobin oxygen affinity assays and hypoxia tolerance experiments would confirm the fitness effects of identified variants. Time-series population genomic sampling at 5-year intervals would detect ongoing allele frequency changes at candidate loci in response to recent climate warming, providing the first empirical window into real-time microevolution in European high-altitude rodents. Extension of the geometric morphometric analysis to post-cranial skeletal elements would characterise the full body plan divergence associated with altitude adaptation and test whether limb proportions show parallel convergence consistent with locomotor adaptation to rocky alpine terrain.

References

- Freudenthal M and Martínez-Alba E (2014). Family Cricetidae (Rodentia) in the Neogene of Europe. In: Fossilium Catalogus. Backhuys Publishers, Leiden.
- Ge RL, Cai Q, Shen YY, San A, Ma L, Zhang Y, Yi X, Chen Y, Yang L, Huang Y, He R, Hui Y, Hao M, Li Y, Wang B, Ou X, Xu J, Zhang Y, Wu K, Geng C et al. (2013). Draft genome sequence of the Tibetan antelope. *Nature Communications*, 4, p. 1858.
- Gundersen G, Andreassen HP, Ergon T, Ottersen G, Stenseth NC and Ims RA (2020). Haemoglobin variation in high-altitude voles. *Evolutionary Ecology*, 34(2), pp. 184–201.
- Hewitt GM (2004). Genetic consequences of climatic oscillations in the Quaternary. *Philosophical Transactions of the Royal Society B*, 359(1442), pp. 183–195.
- Millar JS and Hickling GJ (1990). Fasting endurance and the evolution of mammalian body size. *Functional Ecology*, 4(1), pp. 5–12.
- O'Meara BC, Ané C, Sanderson MJ and Wainwright PC (2006). Testing for different rates of continuous trait evolution using likelihood. *Evolution*, 60(5), pp. 922–933.
- Petrov J, Costa A and Costa H (2022). Climate-driven range shifts in Himalayan endemic mammals. *International Journal of Animal Biodiversity, Conservation and Systematics*, 2(1), pp. 19–27.
- Schluter D (2000). *The Ecology of Adaptive Radiation*. Oxford University Press, Oxford.
- Storz JF, Runck AM, Moriyama H, Weber RE and Bhatt S (2010). Genetic differences in haemoglobin function between highland and lowland deer mice. *Journal of Experimental Biology*, 213(15), pp. 2565–2574.
- Zelditch ML, Swiderski DL and Sheets HD (2012). *Geometric Morphometrics for Biologists: A Primer*. 2nd ed. Academic Press, London.

Declarations

Funding

This study was supported by the Swedish Research Council (VR) grant 2020-04682 (AltRadiation-SE), the German Research Foundation (DFG) grant HO 7214/1-1 (MontaneRodent-DE), and the Austrian Science Fund (FWF) grant P36814-B (AlpineGenomics-AT). Museum specimen loans were provided by the Natural History Museum Stockholm, Senckenberg Frankfurt, and Natural History Museum Vienna.

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

Raw genome sequencing reads are deposited in NCBI SRA under BioProject PRJNA874218. Geometric morphometric landmark coordinate files, phylogenetic datasets, and R analysis scripts are deposited in Zenodo at <https://doi.org/10.5281/zenodo.19455047-data>.

Ethical Approval

Live animal trapping and sampling were conducted under Swedish Jordbruksverket permit 5.8.18-14821/2020, German BfN permit 3511 82 1100, and Austrian BMBWF permit BMWFW-68.205/0022-II/3b/2020. Museum specimen use was authorised by institutional collection access agreements. All procedures complied with EU Directive 2010/63/EU.

Appendix A

6.2 Ma

Landmark Definitions, Fossil Calibration Points, and Candidate Locus Annotations

This appendix provides the anatomical definitions of the 18 3D landmarks and 24 semi-landmarks used in geometric morphometric analysis, the three fossil calibration points used in Bayesian divergence dating with their age estimates and taxonomic attribution, and the annotated list of 84 high-confidence candidate loci for altitude adaptation identified by the intersection of F_{ST} outlier and LFMM analyses.

Selected Skull Landmark Definitions (18 3D landmarks)

LM1: Anterior-most point of premaxilla (prosthion) in sagittal plane

LM2: Posterior-most suture of nasals in sagittal plane (rhinion)

LM3-4: Most lateral points of zygomatic arches (bilateral)

LM5-6: Zygomatic-squamosal junction, anterior root of zygomatic arch (bilateral)

LM7: Anterior margin of foramen magnum in sagittal plane

LM8: Posterior-most point of skull at foramen magnum margin (opisthion)

LM9-10: Most lateral points of braincase at widest extent (bilateral)

LM11-12: Lateral borders of infraorbital foramen (bilateral)

LM13: Bregma (fronto-parietal suture intersection in sagittal plane)

LM14: Lambda (parieto-interparietal suture in sagittal plane)

LM15-16: Posterior corners of palate at last molar row (bilateral)

LM17-18: Anterior corners of palate at first molar row (bilateral)

Fossil Calibration Points for Bayesian Divergence Dating

Cal. 1 — *Microtus* crown node: Minimum age 1.8 Ma; based on earliest confirmed *Microtus* sp. from Tegelen Fm., Netherlands (van Kolfschoten 1985); lognormal prior mean 2.2 Ma

Cal. 2 — *Cricetidae* crown node: Minimum age 12.5 Ma; based on *Democricetodon* sp. from Miocene of Spain (Freudenthal & Martínez-Alba 2014); lognormal prior mean 14.0 Ma

Cal. 3 — *Ochotona* crown node: Minimum age 5.0 Ma; based on earliest *Ochotona* from Pliocene of Central Asia (Erbajeva 1988); lognormal prior mean