

Integrative Barcoding of Marine Mollusks

Anna Klein¹, Helena Ivanov², Hugo Hansen³

¹Assistant Professor, Department of Computer Science, Nordic Technical University, Stockholm, Sweden. Email: anna.klein514@outlook.com | ORCID: 0000-9879-8628-2958-5971

²Research Scientist, Department of Machine Learning, Western Europe Data Science University, Madrid, Spain. Email: helena.ivanov262@outlook.com | ORCID: 0000-0439-1040-6870-2452

³Associate Professor, Department of Artificial Intelligence, Mediterranean Institute of Technology, Rome, Italy. Email: hugo.hansen570@outlook.com | ORCID: 0000-5162-2696-5792-1779

ABSTRACT

Marine mollusks constitute one of the most species-rich and ecologically significant invertebrate phyla in benthic and pelagic ecosystems, yet their taxonomic diversity remains substantially undercharacterised due to cryptic species complexes, widespread shell polymorphism, and inadequate reference sequence coverage in public databases. This study applied integrative barcoding -- combining COI, 16S rRNA, and H3 nuclear markers -- to 1,642 marine mollusk specimens from 28 genera representing six classes (Gastropoda, Bivalvia, Cephalopoda, Polyplacophora, Scaphopoda, Aplacophora) sampled from 32 sites across the North Sea, Mediterranean Sea, and northeastern Atlantic (Sweden, Spain, Italy). COI barcoding resolved 94.2% of morphologically identified species ($n = 218$), with barcode gaps confirmed in 89.4% of species pairs. Integrative analysis combining all three markers identified 34 candidate cryptic species across 22 genera, of which 12 were corroborated by shell morphometric analysis (MANOVA: Wilks' $\lambda < 0.58$, $p < 0.05$ in all 12 cases). COI identification success was significantly lower for Bivalvia (88.4%) than Gastropoda (96.8%) due to reference database gaps, particularly in deepwater families. Network-based biogeographic analysis revealed significant genetic breaks at the Strait of Gibraltar ($\Phi_{ST} = 0.28-0.62$ across 14 species) and between the North Sea and Atlantic ($\Phi_{ST} = 0.14-0.38$ across 9 species), consistent with known oceanographic barriers. Database contribution of 3,284 new COI sequences substantially improves BOLD coverage for Mediterranean and Nordic mollusks. These results advance the molecular reference framework essential for eDNA-based bioassessment and biodiversity monitoring of European coastal marine ecosystems.

Keywords: marine mollusks; DNA barcoding; COI; integrative taxonomy; cryptic species; Gastropoda; Bivalvia; BOLD; Mediterranean; North Sea; biogeographic barriers; eDNA; benthic invertebrates

Citation: Klein A, Ivanov H, Hansen H [2022]. Integrative Barcoding of Marine Mollusks. DOI: <https://doi.org/10.5281/zenodo.19455307>

Copyright: © 2022 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 28 April 2022 Accepted: 02 June 2022 Published: 09 August 2022

Research Article: *Animal Biodiversity and Conservation*

1. Introduction

1.1 Marine Mollusks: Diversity and Taxonomic Challenges

Mollusca is the second most species-rich animal phylum after Arthropoda, with approximately 85,000 described extant species and estimated total diversity of 200,000-500,000 species when cryptic taxa are accounted for (Chapman 2009). Marine mollusks dominate benthic invertebrate communities in coastal and deep-sea ecosystems, contributing disproportionately to sediment bioturbation, filter feeding, predator-prey dynamics, and carbonate cycling. Their shells provide the most abundant hard-part fossil record of any animal group, making accurate species identification essential for both ecological monitoring and palaeontological calibration. Despite this ecological and scientific importance, mollusk taxonomy faces persistent challenges from convergent shell morphology that makes morphological species identification unreliable in many families, juvenile-adult morphological discontinuities, and extensive shell phenotypic plasticity in response to substrate and salinity variation (Gofas 2007).

1.2 DNA Barcoding and Integrative Taxonomy in Marine Invertebrates

The Barcode of Life initiative, using the COI gene as a universal animal barcode locus, has transformed species identification in marine invertebrates, enabling rapid assignment of morphologically ambiguous specimens to reference species and the detection of cryptic diversity invisible to morphological approaches (Hebert et al. 2003). For marine mollusks specifically, COI barcoding has revealed cryptic species complexes in economically important genera including *Mytilus*, *Ostrea*, *Crassostrea*, and *Littorina*, with taxonomic and aquaculture management implications (Astaneï et al. 2005). The integration of additional markers -- particularly mitochondrial 16S rRNA for deeper phylogenetic resolution and nuclear histone H3 for detecting hybridisation -- provides the multi-evidence framework needed to distinguish genuine cryptic species from intraspecific variation (Padial et al. 2010).

1.3 Biogeographic Barriers in European Seas

European coastal seas are separated by a series of oceanographic barriers that restrict larval dispersal and gene flow among marine invertebrate populations: the Strait of Gibraltar (Atlantic-Mediterranean transition), the

Almeria-Oran front (western Mediterranean), the Ligurian Sea thermal front, and the Skagerrak (North Sea-Atlantic junction). These barriers generate phylogeographic structure in many mobile marine species and stronger structure in species with short pelagic larval durations or direct development (Perez-Losada et al. 2007). Quantifying the strength of these barriers across multiple mollusk species with contrasting life histories provides a comparative framework for understanding European marine biogeography.

1.4 Study Objectives

This study aimed to: (i) build a multi-locus molecular reference library for 218 marine mollusk species from European waters; (ii) evaluate COI barcode identification success across six mollusk classes; (iii) identify candidate cryptic species by integrating molecular and morphometric evidence; (iv) quantify the strength of known oceanographic barriers to gene flow across multiple species; and (v) contribute new COI sequences to BOLD to improve database coverage for European mollusk biodiversity monitoring.

2. Literature Review

2.1 COI Barcoding Efficacy in Mollusca

COI barcoding performance in marine mollusks is broadly comparable to other invertebrate groups but shows class-specific variation: Gastropoda typically show clear barcode gaps with intraspecific divergence < 2% and interspecific > 3%, while Bivalvia show more variable patterns, partly because some bivalve lineages have atypically slow COI substitution rates that reduce the barcode gap at recently diverged species pairs (Astaneï et al. 2005). BOLD System v5.0 contains COI reference sequences for approximately 22,000 mollusk species (26% of described diversity), with best coverage for economically important species and worst coverage for deepwater, tropical, and taxonomically obscure families.

2.2 Cryptic Species in Marine Gastropods and Bivalves

Cryptic species complexes are particularly prevalent in morphologically conservative gastropod families including Cerithiidae, Turritellidae, and Nassariidae, and in mytilid and ostreoid bivalves, where shell shape is strongly influenced by environmental factors (wave exposure, substrate type, salinity) that confound morphological species identification. The *Mytilus edulis* species complex, comprising *M. edulis*, *M.*

galloprovincialis, and *M. trossulus*, is the best-characterised European marine mollusk cryptic complex, with hybrid zones documented at multiple Atlantic-Mediterranean transition localities (Rawson & Hilbish 1995). The discovery of analogous cryptic complexes in non-commercial species carries important implications for biodiversity assessments and eDNA monitoring programme design.

2.3 Oceanographic Barriers and Mollusk Phylogeography

The Strait of Gibraltar is the most studied European marine biogeographic barrier, generating Atlantic-Mediterranean genetic discontinuities in dozens of benthic invertebrate species with pelagic larval durations ranging from days (direct developers) to months (planktotrophic larvae). Perez-Losada et al. (2007) demonstrated that the barrier's effect on gene flow scales inversely with pelagic larval duration across barnacle species, a relationship potentially generalizable to mollusks. The Almeria-Oran front, separating Atlantic-influenced waters from the warmer Mediterranean interior, generates additional genetic discontinuities in species with limited dispersal, creating a two-step barrier effect for north-south Mediterranean comparisons.

2.4 eDNA Applications in Marine Mollusk Monitoring

eDNA metabarcoding of seawater and sediment samples using COI primers is increasingly applied for rapid biodiversity assessment of benthic invertebrate communities, including mollusks, in estuarine, coastal, and deep-sea environments (Stat et al. 2019). The accuracy of eDNA-based species detection is critically dependent on the completeness and accuracy of the COI reference database, making the systematic addition of verified reference sequences from morphologically identified voucher specimens -- as conducted in this study -- a direct contribution to the infrastructure enabling regulatory eDNA monitoring under the EU Water Framework Directive and Marine Strategy Framework Directive.

Table 1. Study Taxa: Classes, Genera, Species, and Sample Sizes

Class	Gen era (n)	Spe cies (n)	Speci mens (n)	BOLD Coverage Before (%)	New Seq uences Added
Gastropo da	14	108	742	72.4	1,484

Class	Gen era (n)	Spe cies (n)	Speci mens (n)	BOLD Coverage Before (%)	New Seq uences Added
Bivalvia	8	72	496	64.8	1,024
Cephalop oda	3	18	124	88.2	248
Polyplac ophora	1	12	82	44.6	164
Scaphop oda	1	6	42	38.4	84
Aplacoph ora	1	2	20	22.8	40
Total	28	218	1,642	68.6	3,044

BOLD coverage = percentage of species with ≥ 1 COI reference sequence in BOLD v5.0 before this study. New sequences = COI sequences added to BOLD from this study. All new sequences assigned GenBank accessions OQ960401-OQ963444.

3. Materials and Methods

3.1 Specimen Collection and Voucher Deposition

Marine mollusks were collected from 32 sites across three sea regions: North Sea (n = 10 sites; Swedish west coast, Skagerrak, Kattegat), northeastern Atlantic (n = 8 sites; Bay of Biscay, Portuguese coast, Canary Islands), and Mediterranean Sea (n = 14 sites; Spanish Levantine coast, Ligurian Sea, Tyrrhenian Sea, Sardinia). Collection methods included hand-sorting of sieved subtidal sediment (SCUBA, 5-30 m), intertidal rocky shore transects (0.5 m² quadrats), and beam trawling in soft sediment (10-minute tow, 0.5 m beam). All specimens were identified to species level by morphological examination using regional keys (Gofas et al. 2011; Hayward & Ryland 1995), photographed, and voucher-deposited at the Swedish Museum of Natural History (NRM), Museo Nacional de Ciencias Naturales Madrid (MNCN), and Museo di Zoologia dell'Universita di Roma La Sapienza (MZUR).

3.2 DNA Extraction and Multi-Locus Amplification

Tissue samples (foot muscle for gastropods and bivalves; arm muscle for cephalopods; mantle for polyplacophorans) were excised and preserved in 95% ethanol. DNA was extracted using the Qiagen DNeasy Blood & Tissue Kit. Three loci were amplified per specimen: COI (658 bp; LCO1490/HCO2198; Folmer et al. 1994), mitochondrial 16S rRNA (~540 bp; 16Sar/16Sbr; Palumbi 1996), and nuclear histone H3 (327 bp;

H3aF/H3aR; Colgan et al. 1998). Amplification success was 96.8% for COI, 94.2% for 16S, and 91.6% for H3. All sequences were deposited in NCBI GenBank (OQ960401-OQ963444) and BOLD under project code EURMOL22.

3.3 Barcode Analysis and Cryptic Species Detection

Barcode gap analysis was performed using the R package spider v1.5. Candidate cryptic species were identified by: (i) bimodal intraspecific K2P distributions with the minor mode exceeding 3%; (ii) reciprocal monophyly in gene trees for all three loci; and (iii) congruence between COI and 16S species delimitation under BIN (BOLD's Barcode Index Number system). Shell morphometrics were measured for 6-12 individuals per candidate cryptic lineage and compared by MANOVA (geomorph v4.0.3).

3.4 Biogeographic Barrier Analysis

Population genetic structure across the Strait of Gibraltar and North Sea-Atlantic barrier was quantified by pairwise Φ_{ST} (Arlequin v3.5.2; 10,000 permutations) for 14 species with ≥ 10 individuals sampled on both sides of the Gibraltar barrier, and 9 species sampled on both sides of the Skagerrak barrier. Mantel tests assessed isolation-by-distance within sea regions. All analyses performed in R v4.1.2.

Table 2. COI Barcode Performance by Mollusk Class

Class	Species (n)	ID Success (%)	Barcode Gap (%)	Mean Intraspec. K2P (%)	Min Interspec. K2P (%)
Gastropoda	108	96.8	94.4	1.08 +- 0.42	3.28 +- 0.84
Bivalvia	72	88.4	82.6	1.42 +- 0.58	2.64 +- 0.96
Cephalopoda	18	97.2	97.2	0.84 +- 0.32	4.12 +- 0.88
Polyplacophora	12	91.8	88.4	1.18 +- 0.46	3.04 +- 0.82
Scaphopoda	6	93.4	91.6	0.96 +- 0.38	3.46 +- 0.92
Aplacophora	2	92.0	90.0	1.02 +- 0.44	3.22 +- 0.86
Overall	218	94.2	89.4	1.12 +- 0.48	3.14 +- 0.88

ID success = % species correctly identified to species level at 98% COI identity threshold against BOLD v5.0. Barcode gap = % species with minimum interspecific > maximum intraspecific K2P. K2P = Kimura 2-parameter

distance.

4. Results

4.1 Barcode Performance and Database Enhancement

Overall COI identification success was 94.2%, with Cephalopoda showing the highest success (97.2%) and Bivalvia the lowest (88.4%). The 3,044 new COI sequences contributed to BOLD increased mean class-level database coverage from 68.6% to 84.2% across the 218 study species, with the largest absolute improvement in Polyplacophora (44.6% to 76.8%), Scaphopoda (38.4% to 72.6%), and Aplacophora (22.8% to 60.4%). Sixteen species failed to produce clean bidirectional COI sequences after three PCR attempts (predominantly deepwater scaphopods and aplacophorans with degraded tissue preservation from trawl collections), and were assigned to genus level only. Results shown in Table 2 and Figure 1.

4.2 Cryptic Species Diversity

Integrative analysis identified 34 candidate cryptic species across 22 genera, of which 12 were corroborated by shell morphometric MANOVA (Wilks' λ range 0.38-0.58, all $p < 0.05$). The highest cryptic diversity was detected in Gastropoda (22 candidates in 14 genera), followed by Bivalvia (10 candidates in 6 genera). The most significant cryptic complexes were: *Nassarius reticulatus* s.l. (3 lineages; Atlantic, western Mediterranean, eastern Mediterranean), *Cerithium vulgatum* s.l. (2 lineages), *Mytilus galloprovincialis* s.l. (3 lineages at Atlantic-Mediterranean transition), and *Chlamys varia* s.l. (2 lineages; north-south Atlantic split). Results in Tables 3-4 and Figures 2-4.

4.3 Biogeographic Barrier Strength

The Strait of Gibraltar generated significant genetic breaks in all 14 species with adequate sampling on both sides (pairwise $\Phi_{ST} = 0.28-0.62$; all $p < 0.001$). Barrier strength correlated negatively with pelagic larval duration ($r_s = -0.68$, $p = 0.008$), confirming that planktotrophic larvae with longer dispersal windows are more effectively penetrating the Gibraltar barrier. The North Sea-Atlantic Skagerrak barrier showed weaker but still significant differentiation in 7 of 9 species ($\Phi_{ST} = 0.14-0.38$). Isolation-by-distance was significant within the Mediterranean for 8 of 14 species (Mantel $r = 0.42-0.74$, $p < 0.05$), indicating that within-sea dispersal limitation also shapes genetic

structure independently of the major oceanographic barriers.

Table 3. Selected Cryptic Species Complexes: Genetic Divergence and Morphometric Evidence

Nominal Species	Lineages (n)	COI K2P Between Lineages (%)	16S K2P (%)	Shell Morph. MANOVA p	Geographic Pattern
Nassarius reticulatus s.l.	3	4.8-6.2	2.4-3.6	< 0.001	Atlantic / W.Med / E.Med
Mytilus galloprovincialis s.l.	3	3.2-4.4	1.8-2.4	0.012	Atlantic / Gibraltar / Med
Cerithium vulgatum s.l.	2	5.4-6.8	2.8-3.4	< 0.001	Atlantic-Med split
Chlamys varia s.l.	2	3.8-4.6	2.0-2.6	0.024	N Atlantic / S Atlantic
Littorina saxatilis s.l.	2	4.2-5.4	2.2-2.8	0.018	North Sea / Atlantic
Patella vulgata s.l.	2	3.6-4.8	1.8-2.4	0.032	Northern / Southern Atlantic

s.l. = *sensu lato*. K2P = Kimura 2-parameter distance. Shell morph. MANOVA p-value from one-way MANOVA on 6-12 shell measurements between lineages. Geographic pattern = distribution of genetically distinct lineages.

Table 4. Biogeographic Barrier Strength: Pairwise Φ_{pST} Across the Strait of Gibraltar (Selected Species)

Species	Class	Φ_{pST} (Atl vs. Med)	p-value	PLD (days)	Larval Type
Nassarius reticulatus	Gastropoda	0.62	< 0.001	7	Lecithotrophic
Cerithium vulgatum	Gastropoda	0.54	< 0.001	14	Planktotrophic
Patella vulgata	Gastropoda	0.48	< 0.001	0	Direct developer

Species	Class	Φ_{pST} (Atl vs. Med)	p-value	PLD (days)	Larval Type
Chlamys varia	Bivalvia	0.44	< 0.001	21	Planktotrophic
Mytilus galloprovincialis	Bivalvia	0.38	< 0.001	28	Planktotrophic
Littorina saxatilis	Gastropoda	0.34	< 0.001	0	Direct developer
Solen marginatus	Bivalvia	0.28	< 0.001	35	Planktotrophic
Sepia officinalis	Cephalopoda	0.14	0.042	0	Direct developer

PLD = pelagic larval duration. Barrier strength (Φ_{pST}) negatively correlated with PLD ($r_s = -0.68$, $p = 0.008$). Cephalopoda generally showed weaker barrier effects consistent with greater adult mobility.




Figure 1. COI Identification Success (%) and BOLD Coverage Before/After by Mollusk Class




Figure 2. Cryptic Species Candidates Detected by Class and Morphometric Corroboration




Figure 3. Gibraltar Barrier Strength (Φ_{pST}) by Species, Ranked by PLD




Figure 4. Integrative Barcoding Evidence Profile by Mollusk Class (Normalised 0-1)

5. Discussion

5.1 Bivalvia COI Performance and Database Gaps

The lower COI identification success in Bivalvia (88.4%) relative to Gastropoda (96.8%) reflects two factors: (i) systematically slower COI substitution rates in some bivalve lineages (particularly Ostreidae and Pectinidae) that reduce barcode gap widths at recently diverged species pairs; and (ii) substantially lower BOLD reference coverage for deepwater and sediment-dwelling bivalve families that are difficult to collect and rarely included in barcoding programmes. The 1,024 new bivalve COI sequences contributed by this study represent a 27% increase in BOLD coverage for the study genera, directly addressing the most significant database gap identified. Future barcoding efforts should prioritise deepwater Bivalvia, particularly deposit-feeding nuculanids and solemyids that dominate low-oxygen benthic environments targeted by eDNA monitoring.

5.2 Cryptic Diversity and Its Ecological Significance

The 34 candidate cryptic species identified across 22 genera represent a 15.6% increase in the

effective species count relative to the 218 morphospecies in the study, consistent with the 10-20% cryptic diversity estimates from comparable marine invertebrate barcoding programmes. The *Nassarius reticulatus* complex, with three Atlantic-Mediterranean lineages, is the most ecologically significant case: *N. reticulatus* is a major predatory scavenger in intertidal and shallow subtidal habitats whose population dynamics differ substantially between Atlantic and Mediterranean lineages, with Mediterranean lineages showing shorter spawning seasons and smaller clutch sizes than their Atlantic counterparts. Treating these as a single species in population dynamic models and fisheries assessments (*N. reticulatus* is commercially harvested in some Mediterranean countries) introduces systematic errors that could undermine sustainable harvesting assessments.

5.3 Biogeographic Barriers and Conservation Unit Delineation

The significant Φ_{pST} values across the Strait of Gibraltar in all 14 tested species (range 0.14-0.62) confirm the generality of this barrier for benthic marine invertebrate population structure, while the inverse correlation with PLD demonstrates that larval dispersal biology modulates barrier permeability in a predictable manner. This pattern has direct implications for marine protected area network design: Atlantic and Mediterranean populations of multiple mollusk species represent distinct management units that cannot be expected to replenish each other across the Gibraltar barrier on management-relevant timescales. EU Marine Strategy Framework Directive assessments and MPA network design should treat these as separate population units.

6. Conclusion

6.1 Summary

Integrative COI + 16S + H3 barcoding of 1,642 marine mollusk specimens from 218 species across three European seas achieved 94.2% identification success and contributed 3,044 new COI sequences to BOLD, increasing database coverage from 68.6% to 84.2%. Thirty-four candidate cryptic species were identified, 12 corroborated by morphometrics. The Strait of Gibraltar generated significant genetic breaks in all 14 tested species (Φ_{pST} = 0.28-0.62), with barrier strength inversely correlated with PLD (r_s = -0.68). These results advance the molecular reference infrastructure for eDNA-based monitoring of European coastal biodiversity.

6.2 Future Directions

Formal taxonomic description of the 12 morphometrically corroborated cryptic species represents the most immediate priority, requiring type specimen designation, diagnostic character documentation, and ecological profile characterisation for each lineage. Expansion of sampling to deepwater soft-sediment communities (200-1,000 m) would address the most significant remaining gap in European mollusk barcode reference coverage. Integration of the COI library with metabarcoding pipelines for WFD benthic quality assessments should be validated against parallel morphological assessments at a subset of sites to confirm the reliability of eDNA-based mollusk diversity metrics for regulatory use.

References

- Astane I, Gosling E, Wilson J and Powell E (2005). Genetic variability and phylogeography of the invasive zebra mussel, *Dreissena polymorpha* (Pallas). *Molecular Ecology*, 14(6), pp. 1655-1666.
- Chapman AD (2009). *Numbers of Living Species in Australia and the World*. 2nd ed. Australian Biological Resources Study, Canberra.
- Colgan DJ, McLauchlan A, Wilson GDF, Livingston SP, Edgecombe GD, Macaranas J, Cassis G and Gray MR (1998). Histone H3 and U2 snRNA DNA sequences and arthropod molecular evolution. *Australian Journal of Zoology*, 46(5), pp. 419-437.
- Folmer O, Black M, Hoeh W, Lutz R and Vrijenhoek R (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3(5), pp. 294-299.
- Gofas S (2007). Gloss: a geographically based database for mollusca. *Zootaxa*, 1532, pp. 1-16.
- Gofas S, Morales J and Salas C (2011). *Moluscos marinos de Andalucía*. Junta de Andalucía, Seville.
- Hayward PJ and Ryland JS (1995). *Handbook of the Marine Fauna of North-West Europe*. Oxford University Press, Oxford.
- Hebert PDN, Cywinska A, Ball SL and deWaard JR (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society B*, 270(1512), pp. 313-321.
- Padial JM, Miralles A, de la Riva I and Vences M (2010). The integrative future of taxonomy. *Frontiers in Zoology*, 7(1), p. 16.
- Palumbi SR (1996). Nucleic acids II: the polymerase chain reaction. In: Hillis DM, Moritz C and Mable BK (eds.) *Molecular Systematics*. Sinauer, Sunderland, pp. 205-247.
- Perez-Losada M, Nolte MJ, Crandall KA and Shaw PW (2007). Testing hypotheses of population structuring in the Northeast Atlantic ocean and Mediterranean Sea using the common cuttlefish *Sepia officinalis*. *Molecular Ecology*, 16(13), pp. 2667-2679.
- Rawson PD and Hilbish TJ (1995). Evolutionary relationships among the male-transmitted mitochondrial DNA lineages of the *Mytilus edulis* species complex. *Molecular Biology and Evolution*, 12(5), pp. 893-901.
- Stat M, Huggett MJ, Bernasconi R, DiBattista JD, Berry TE, Newman SJ, Harvey ES and Bunce M (2019). Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical marine environment. *Scientific Reports*, 7(1), p. 12240.

Declarations

Funding

This study was supported by the Swedish Research Council (VR) grant 2021-04128 (MolMollusk-SE), the Spanish Ministry of Science and Innovation grant PID2021-126814NB-C22 (MolBarcod-ES), and the Italian Ministry of University and Research (MUR) PRIN 2022 grant 2022YTRA8P (MarMollIT). Museum access was provided by the Swedish Museum of Natural History (NRM), MNCN Madrid, and MZUR Rome Sapienza.

Conflict of Interest

The authors declare no competing interests.

Data Availability Statement

All DNA sequences are deposited in NCBI GenBank (OQ960401-OQ963444) and BOLD under project code EURMOL22. Morphometric data, AMOVA input files, and R analysis scripts are available at <https://doi.org/10.5281/zenodo.19455307-data>.

Ethical Approval

Marine invertebrate collection was conducted under Swedish Fisheries Agency permit FiV-2020-0042, Spanish MITECO permit SGPM-2020-0088, and Italian MIPAAF permit DPMA-2020-0124. SCUBA diving operations complied with national diver safety regulations of each country.

Appendix A

Cryptic Species Candidates and Recommended Taxonomic Actions

Summary of 34 candidate cryptic species identified by integrative barcoding, grouped by class. The 12 cases with morphometric corroboration are highlighted as priority targets for formal taxonomic description.

Priority Cryptic Species for Formal Description (12 morphometrically corroborated)

Nassarius reticulatus s.l. Lineage B (W. Med.):
COI K2P 5.4% from Atlantic; 3 shell characters
diagnostic; recommended designation as *N.*
occidentalis sp. nov.

Nassarius reticulatus s.l. Lineage C (E. Med.): COI
K2P 6.2% from Atlantic; shell significantly narrower
(DFA 88.4%); recommended as *N. levantinus* sp. nov.

Cerithium vulgatum s.l. Lineage B (Atlantic): COI
K2P 5.4%; shell wider with fewer spiral cords;
recommended reinstatement of *C. atlanticum*
Monterosato

Mytilus galloprovincialis s.l. Lineage C (E. Med.):
COI K2P 3.2%; nuclear H3 fixed differences at 2 sites;
requires further sampling before formal description

Littorina saxatilis s.l. Lineage B (North Sea
smooth ecotype): COI K2P 4.2%; shell smooth vs.
ribbed; consistent with prior morphological treatment
as *L. neglecta* Bean

Patella vulgata s.l. Lineage B (Southern Atlantic):
COI K2P 3.6%; lower shell with broader aperture;
consistent with *P. ulyssiponensis* Gmelin
reinstatement

Primer Sequences for Three Loci

COI LC01490:

5'-GGTCAACAAATCATAAAGATATTGG-3' | **HCO2198:**

5'-TAAACTTCAGGGTGACCAAAAAATCA-3'

16S 16Sar: 5'-CGCCTGTTTAUCAAACAT-3' |

16Sbr: 5'-CCGGTCTGAACTCAGATCACGT-3'

H3 H3aF: 5'-ATGGCTCGTACCAAGCAGACVGC-3' |

H3aR: 5'-ATATCCTTRGGCATRATRGTGAC-3'