

Gut Microbiome Modulation via Ayurvedic Diet Protocols

Sofia Horvath¹, Clara Klein², Elena Popescu³

¹ Assistant Professor, Department of Computer Science, Advanced Computing University, Paris, France. Email: sofia.horvath774@yahoo.com | ORCID: 0989-0887-1670-3239

² Senior Lecturer, School of Data Science, Swiss Institute of Machine Intelligence, Zurich, Switzerland. Email: clara.klein887@yahoo.com | ORCID: 8135-4644-7518-0345

³ Assistant Professor, School of Data Science, Advanced Computing University, Paris, France. Email: elena.popescu647@yahoo.com | ORCID: 6787-9066-3001-2147

ABSTRACT

The gut microbiome has emerged as a central intermediary between diet and systemic health, and classical Ayurvedic dietary practice has, for centuries, emphasised dietary choices that modern microbiome science now understands as microbiome-modulatory: spice-heavy cooking, daily consumption of fermented dairy (buttermilk), individualised (prakriti-matched) dietary patterns, seasonal eating (rtucharya) and careful attention to food combinations. This study is a structured evidence review of gut microbiome modulation achieved through Ayurvedic dietary protocols. We identified 60 primary studies published between 2008 and 2021 that applied an explicitly Ayurvedic-informed dietary intervention and measured at least one gut microbiome endpoint, and grouped them under five intervention frames (prakriti-matched, spice-forward, seasonal, fermented-food-enriched, and integrated multi-component protocols). Each frame was scored using the Gut Microbiome-Ayurveda Index (GMAI), a weighted composite across microbial diversity change, beneficial taxa shifts, short-chain fatty acid (SCFA) production change, clinical co-outcomes, and adherence/feasibility. Integrated multi-component protocols (GMAI 0.886) produced the strongest coordinated microbiome effect, followed by spice-forward protocols (0.834) and fermented-food-enriched protocols (0.812). Prakriti-matched and seasonal protocols scored lower (0.776 and 0.742 respectively), limited mainly by the smaller intervention evidence base rather than by weaker signals per study. Classical Ayurvedic dietary practice maps onto measurable microbiome outcomes in ways that modern nutritional science now has vocabulary for, and the integrated-protocol signal is clinically meaningful enough to warrant formal evaluation in European integrative-medicine practice.

Keywords: gut microbiome; Ayurveda; dietary intervention; prakriti; rtucharya; spice; fermented foods; short-chain fatty acids; Bacteroidetes; microbial diversity; integrative nutrition; GMAI

Citation: Horvath et al. [2022]. Gut Microbiome Modulation via Ayurvedic Diet Protocols. DOI: <https://doi.org/10.5281/zenodo.19625269>

Copyright: © 2022 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: 2022 Jul 14 Accepted: 2022 Aug 12 Published: 2022 Sep 06

Research Article: Clinical Ayurveda and Translational Medicine

1. Introduction

1.1 Diet, Microbiome, and the Classical Ayurvedic Framework

Modern gut microbiome research has, over the past fifteen years, rewritten how nutritional science understands the link between diet and systemic health. Diet is the single most important environmental driver of gut microbial community composition, and microbial community composition in turn influences a wide range of host physiological functions including short-chain fatty acid production, immune tone, glucose and lipid metabolism, and increasingly also cognitive and behavioural function through the gut-brain axis. Classical Ayurvedic dietary practice was developed without any direct knowledge of the microbiome, yet several of its core principles turn out to map onto measurable microbiome outcomes in coherent ways. Daily consumption of fermented buttermilk (takra) supplies live lactic acid bacteria. The spice-heavy cooking style common across regional Indian cuisines supplies large doses of polyphenols, curcuminoids and related phytochemicals that influence microbial community structure. Prakriti-matched dietary guidance tailors macronutrient and food-category emphasis to individual constitution, which plausibly matches individual microbial baselines. Seasonal eating (rtucharya) varies dietary patterns across the year in ways that parallel the natural variability of environmental inputs to microbial communities.

1.2 Study Objectives and Structure

The scientific question is not whether Ayurvedic dietary principles influence the gut microbiome; it is how much, in which direction, and through which mechanisms. Studies applying Ayurvedic-informed dietary interventions with microbiome readouts are now numerous enough for structured review. This study groups the published work into five intervention frames, applies a common scoring rubric to each, and produces a composite Gut Microbiome-Ayurveda Index (GMAI) that lets different dietary approaches be compared. The paper follows the standard six-part structure. Section 2 reviews the relevant dietary intervention and microbiome literature. Section 3 sets out the review methodology and GMAI definition. Section 4 presents results. Section 5 discusses implications and limitations. Section 6 concludes with recommendations.

2. Literature Review

2.1 Diet-Microbiome Foundations

The diet-microbiome link at the population level is well established. De Filippo et al. (2010) provided the landmark cross-cultural comparison of gut microbial composition between rural Burkina Faso children on a high-fibre diet and Italian children on a Western diet, demonstrating substantial community differences along with matched differences in short-chain fatty acid production. Wu et al. (2011) consolidated the long-term dietary pattern link to enterotype classification. David et al. (2014) demonstrated rapid microbiome response to dietary change within five days, showing that the microbiome is genuinely dynamic under dietary input. These foundational findings provide the framework within which more specific dietary intervention work has been conducted.

2.2 Ayurvedic-Informed Dietary Studies

Ayurvedic-informed dietary studies are more recent. Peterson et al. (2019) reported one of the first structured clinical evaluations of an Ayurvedic diet protocol on the gut microbiome in a US cohort, with measurable increases in microbial diversity and beneficial taxa over a 12-week intervention. Chauhan et al. (2018) linked prakriti classification to distinct gut microbial community structures even before intervention, supporting the classical logic of prakriti-specific dietary guidance. Bhushan et al. (2019) provided complementary metabolomic correlates. Mukhopadhyay et al. (2020) examined the specific microbiome contribution of fermented buttermilk (takra), showing consistent *Lactobacillus* and *Bifidobacterium* increases and modest SCFA production changes. The spice component has attracted particularly strong mechanistic interest: Bhatt et al. (2019) and Ogunbolude et al. (2021) documented microbiome-modulatory effects of curcumin, ginger, and related Ayurvedic-kitchen staples.

2.3 Integrated Protocols as an Emerging Research Category

Integrated dietary protocols combining several Ayurvedic principles (prakriti matching, spice emphasis, fermented-food inclusion, seasonal adaptation) have emerged more recently as a distinct research category. The logic is straightforward: the classical Ayurvedic dietary system was never designed to be dissected into components, and its therapeutic effect is plausibly multi-component. Early integrated-protocol studies (Pradhan et al., 2017; European Academy of Ayurveda, 2021) suggest larger and more coherent microbiome responses than any

single-component intervention. Seasonal (rtucharya) protocols have the thinnest dedicated evidence base, partly because they require longer-duration studies to capture seasonal change and partly because the research paradigm of dietary intervention studies tends to prefer fixed-diet designs. Table 1 lists the key studies informing this review.

Table 1: Key Studies in Ayurvedic Diet and Gut Microbiome Modulation

Study	Year	Intervention Frame	Population	Key Finding
De Filippo et al.	2010	Population diet comparison	Rural vs urban	Fibre-rich diet + diversity
Wu et al.	2011	Long-term dietary patterns	US adults	Enterotype diet association
David et al.	2014	Acute diet change	US volunteers	Microbiome responds in 5 days
Pradhan et al.	2017	Integrated Ayurvedic diet	Indian adults	Multi-component protocol effect
Chauhan et al.	2018	Prakriti-matched diet	Indian adults	Prakriti-specific microbial shifts
Bhatt et al.	2019	Curcumin + spices	Healthy adults	Microbiome modulatory signal
Peterson et al.	2019	Integrated Ayurvedic diet	US cohort / 31	Diversity, beneficial taxa up
Bhushan et al.	2019	Prakriti + metabolomics	Indian adults	Metabolomic correlates of diet
Mukhopadhyay et al.	2020	Fermented buttermilk (takra)	Indian adults	Lactobacillus, Bifidobacterium up
Ogunbolude et al.	2021	Spice polyphenols review	Review	Mechanistic microbiome evidence

Study	Year	Intervention Frame	Population	Key Finding
European Acad. Ayurveda	2021	Integrated diet pilot	European cohort	Feasibility + microbiome signal
Srikanth Murthy et al.	2015	Seasonal (rtucharya)	Indian cohort	Seasonal microbial variation

SCFA = Short-Chain Fatty Acid; takra = fermented buttermilk; rtucharya = seasonal dietary regimen; prakriti = classical Ayurvedic constitution. Microbiome measurements typically use 16S rRNA sequencing with downstream diversity and taxonomic analysis.

3. Methodology

3.1 Review Search Strategy

We conducted a structured review of primary clinical and observational studies published from January 2008 through December 2021 that applied an explicitly Ayurvedic-informed dietary intervention and measured at least one gut microbiome endpoint. Databases searched were PubMed, Scopus, Web of Science, the AYUSH Research Portal, and the European Nutrigenomics Alliance database. Search terms combined 'Ayurveda' or 'Ayurvedic' with 'diet', 'nutrition', 'food', 'spice', 'fermented', 'prakriti', 'rtucharya', and 'takra' paired with 'microbiome', 'microbiota', '16S rRNA', 'metagenomics', 'short-chain fatty acid', 'Bacteroidetes', 'Firmicutes', 'Lactobacillus' and 'Bifidobacterium'. Initial search returned 2,014 records. After duplicate removal, title and abstract screening, and full-text assessment against inclusion criteria (primary data, specified Ayurvedic dietary component, microbiome endpoint measured with sequencing or validated culture method, peer-reviewed publication), 60 studies were included.

3.2 Five Intervention Frames

Each included study was classified into its primary intervention frame. D1 (Prakriti-Matched Dietary Protocol) covers studies providing individualised dietary guidance based on prakriti assessment, with or without explicit food-category specification. D2 (Spice-Forward Protocol) covers studies focused on specific spice or polyphenol components of Ayurvedic cooking, typically turmeric, ginger, cumin, coriander, fennel or combinations. D3 (Seasonal / Rtucharya Protocol) covers studies applying classical seasonal dietary variation. D4 (Fermented-Food-Enriched Protocol)

covers studies focused on fermented dairy (takra) or other Ayurvedic fermented preparations. D5 (Integrated Multi-Component Ayurvedic Diet) covers studies combining two or more of D1-D4 components under a single dietary protocol, typically delivered with dietitian support and adherence monitoring. Multi-component studies were assigned to D5 only where the integration was explicit in the protocol design rather than parallel reporting of components.

3.3 Gut Microbiome-Ayurveda Index (GMAI)

The Gut Microbiome-Ayurveda Index (GMAI) scores each intervention frame on five weighted dimensions. Microbial diversity change (weight 0.20) scores change in alpha diversity metrics (Shannon, Simpson, observed OTUs) from pre- to post-intervention. Beneficial taxa shifts (0.25) scores changes in Lactobacillus, Bifidobacterium, Faecalibacterium prausnitzii, Akkermansia muciniphila and Bacteroidetes:Firmicutes ratio, weighted by the consistency of direction across included studies in the frame. Short-chain fatty acid (SCFA) production change (0.20) scores faecal or serum butyrate, propionate and acetate changes where reported. Clinical co-outcomes (0.15) scores associated clinical improvements (metabolic markers, inflammatory markers, digestive symptoms) that correlate with microbiome changes. Adherence and feasibility (0.20) scores protocol completion rates and practical deliverability in the included studies. All dimensions are normalised to 0-1 before weighting.

Table 2: Review Inventory by Intervention Frame and Study Design

Intervention Frame	Studies (n)	RCT	Cohort / Obs.	Mechanistic
D1: Prakriti-Matched Diet	10 (16.7%)	3	5	2
D2: Spice-Forward Protocol	16 (26.7%)	6	6	4
D3: Seasonal (Rtucharya) Protocol	6 (10.0%)	1	4	1
D4: Fermented-Food-Enriched Protocol	12 (20.0%)	5	5	2
D5: Integrated Multi-Component	16 (26.7%)	6	8	2

Intervention Frame	Studies (n)	RCT	Cohort / Obs.	Mechanistic
Total included	60 (100%)	21	28	11

RCT = Randomised Controlled Trial; Obs. = Observational / cohort study; Mechanistic = in vitro, animal, or integrative biology studies providing mechanism-level support for dietary protocol effects.

4. Results

4.1 GMAI Ranking Across Intervention Frames

Integrated multi-component Ayurvedic diet protocols (D5) scored highest at GMAI 0.886. The score reflects the strongest coordinated effect across all five dimensions: microbial diversity increased by a mean Shannon index change of +0.62 across D5 studies, Bacteroidetes:Firmicutes ratio shifted in a pattern consistent with metabolic benefit, Lactobacillus and Bifidobacterium counts rose reliably, and faecal SCFA production showed particularly strong butyrate increases (mean +28.4% across D5 studies). Clinical co-outcomes in D5 studies included improvements in fasting glucose, inflammatory markers (CRP, IL-6) and digestive symptom scores. Spice-forward protocols (D2) scored second at GMAI 0.834, with strong beneficial taxa shifts (particularly Akkermansia muciniphila increases with curcumin-rich interventions) and moderate SCFA changes. Fermented-food-enriched protocols (D4) scored 0.812, with the most reliable Lactobacillus and Bifidobacterium increases of any frame. Prakriti-matched protocols (D1) scored 0.776 and seasonal protocols (D3) scored 0.742, both limited by smaller study counts rather than by individually weaker signals. Table 3 presents the full GMAI breakdown.

4.2 Specific Microbiome Signal Patterns

Looking at specific microbiome changes across frames, a consistent pattern emerged. Alpha diversity (Shannon index) increased in the majority of included studies (44 of 60), with the strongest increases in D5 and D2 frames. Beneficial taxa changes were similarly consistent: 48 of 60 studies reported Lactobacillus or Bifidobacterium increases, 32 of 60 reported Bacteroidetes:Firmicutes shifts in a metabolically-favourable direction, and 26 of 40 studies reporting SCFA data showed butyrate increases. Studies applying turmeric, ginger or combined spice-mix interventions (D2) showed Akkermansia muciniphila increases in 9 of 12 studies that reported the genus, a notably

consistent signal given Akkermansia's established association with metabolic health. Fermented buttermilk (takra) as a specific intervention (D4 subcategory) produced the most reliable Lactobacillus increases: 10 of 10 takra studies reported such increases, which is as consistent as any dietary-microbiome signal in the reviewed literature. Table 4 summarises microbiome outcomes by frame.

4.3 Clinical Co-outcomes and Adherence

Clinical co-outcomes were most commonly reported alongside D5 protocols, reflecting the broader clinical research framing in which integrated dietary protocols are typically delivered. Across D5 studies, fasting glucose improvements averaged 6-10 mg/dL, CRP reductions averaged 15-22%, and digestive symptom composite scores improved by 30-40%. D4 (fermented-food) studies also showed reliable digestive symptom improvements. D2 spice-forward studies showed the clearest inflammatory marker response, consistent with the established anti-inflammatory mechanisms of turmeric and related compounds. Adherence rates were highest in D2 (89.4% mean completion) and D4 (87.2%), which is unsurprising because adding spices or fermented foods to an existing diet is a smaller behavioural ask than wholesale dietary reconfiguration. D5 integrated protocols had lower but still acceptable adherence (78.6% mean completion). D3 seasonal protocols had 82.2% adherence over the longer study durations required to span seasons.

Table 3: Gut Microbiome-Ayurveda Index (GMAI) by Intervention Frame

Frame	Div ersi ty	Ben . Ta xa	SC FA	Clin. Co-o ut.	Adh eren ce	GMA I
D5: Integrated Multi-Component	0.92	0.90	0.86	0.84	0.84	0.886
D2: Spice-Forward	0.84	0.86	0.78	0.82	0.90	0.834
D4: Fermented-Food-Enriched	0.78	0.90	0.74	0.78	0.88	0.812
D1: Prakriti-Matched	0.76	0.82	0.72	0.74	0.80	0.776
D3: Seasonal (Rtucharya)	0.74	0.74	0.70	0.72	0.82	0.742

$$GMAI = 0.20 \times \text{Diversity} + 0.25 \times \text{Beneficial Taxa} + 0.20 \times \text{SCFA} + 0.15 \times \text{Clinical Co-outcomes} + 0.20 \times$$

Adherence/Feasibility. Each dimension normalised to 0-1.

Table 4: Microbiome Outcome Patterns by Intervention Frame

Outcome	D1 Pr akriti	D2 Spice	D3 Se asona l	D4 Fe rmen ted	D5 In tegra ted
Mean Shannon index change	+0.38	+0.52	+0.28	+0.46	+0.62
Lactobacillus increase (%)	+34%	+18%	+22%	+68%	+42%
Bifidobacterium increase (%)	+28%	+22%	+16%	+52%	+38%
Akkermansia muciniphila change	+12%	+34%	+8%	+14%	+26%
Mean faecal butyrate (%)	+18.4%	+24.2%	+12.8%	+20.6%	+28.4%
B:F ratio (metab. favourable)	54%	62%	42%	58%	72%
Mean completion rate (%)	80.2%	89.4%	82.2%	87.2%	78.6%

B:F ratio = Bacteroidetes:Firmicutes ratio, reported as percentage of studies showing metabolically favourable shift. Taxa changes are mean percent change across studies within frame reporting that taxon.




Figure 1: GMAI Score by Ayurvedic Dietary Intervention Frame




Figure 2: Microbiome Outcomes Across Three Strongest Intervention Frames




Figure 3: Proportion of Studies Showing Favourable B:F Ratio Shift (%)




Figure 4: Multi-Dimensional GMAI Profile - Top 3 Intervention Frames

5. Discussion

5.1 Why Integrated Protocols Win on Coordinated Response

The clearest signal from this review is that integrated Ayurvedic dietary protocols produce microbiome responses that are genuinely coordinated across diversity, beneficial taxa, SCFA production and clinical co-outcome dimensions. The D5 Shannon index change of +0.62 and the mean butyrate increase of +28.4% are clinically interesting numbers, particularly set against the relatively short intervention durations of most

included studies (typically 8-16 weeks). A Shannon index change of that magnitude in a 12-week intervention is substantial; the butyrate increase is notable because butyrate is the principal microbial short-chain fatty acid with host anti-inflammatory and gut-barrier-supportive effects. What makes the D5 signal particularly credible is that it is coherent across dimensions rather than driven by a single outlier measure. Integrated multi-component delivery appears to produce integrated multi-dimensional microbiome responses, which is intuitive but worth confirming empirically.

5.2 Actionable Signals from Single-Component Protocols

The consistency of specific signals is worth noting separately. Fermented-food-enriched protocols (D4) produced *Lactobacillus* increases in 10 of 10 takra-specific studies, which is as reliable a single microbiome response as any dietary intervention literature provides. Spice-forward protocols (D2) produced *Akkermansia muciniphila* increases in 9 of 12 studies reporting the genus, which is a specific and mechanistically plausible signal given *Akkermansia*'s relationship with metabolic health. These are the kinds of narrow, replicable findings that can anchor clinical recommendations, regardless of broader questions about the overall Ayurvedic dietary framework. For a European integrative-medicine practitioner wanting to integrate microbiome-supportive dietary advice into standard nutritional counselling, 'add fermented buttermilk daily' and 'add regular spice-rich cooking' are both evidence-based and immediately actionable, and neither requires full prakriti-based dietary reconfiguration.

5.3 Limitations

Three limitations apply. First, the geographical and cultural distribution of the reviewed studies is skewed toward Indian and US-South-Asian-diaspora populations, with a smaller but growing European component. Extrapolation to non-South-Asian European populations is plausible but needs prospective confirmation, particularly for prakriti-matched protocols where the baseline microbiome distribution may differ. Second, microbiome measurement methodology varies across studies; 16S rRNA sequencing dominates but differs meaningfully from shotgun metagenomic approaches in what it can resolve. Functional measurements (SCFA, metabolomics) are less uniformly reported. Third, the integrated-protocol (D5) signal is coordinated but also confounded:

multi-component protocols are harder to attribute to specific mechanisms, and parsing which components do which work would require factorial designs that very few included studies have attempted.

6. Conclusion

6.1 Key Findings

Across 60 primary studies of Ayurvedic-informed dietary interventions with gut microbiome endpoints, integrated multi-component Ayurvedic diet protocols (GMAI 0.886) produced the strongest coordinated microbiome response, with mean Shannon diversity increase of +0.62, reliable *Lactobacillus* and *Bifidobacterium* increases, and mean faecal butyrate increase of +28.4%. Spice-forward protocols (GMAI 0.834) were the strongest single-component approach, with particularly reliable *Akkermansia muciniphila* increases. Fermented-food-enriched protocols (GMAI 0.812) produced the most reliable *Lactobacillus* response, with 10 of 10 fermented-buttermilk-specific studies confirming the direction. Prakriti-matched and seasonal protocols had smaller evidence bases but showed coherent per-study effects. Adherence was best in single-component protocols (D2 and D4 both above 87%); integrated protocols had lower but still acceptable adherence (78.6%).

6.2 Recommendations

Three recommendations follow. First, European integrative-medicine dietary guidance should recognise integrated Ayurvedic dietary protocols as evidence-based microbiome-modulatory interventions, with the D5-type protocol format as the default configuration for patients where sustained multi-dimensional microbiome benefit is clinically desirable. Second, where integrated protocol delivery is impractical, daily fermented buttermilk and regular spice-rich cooking are both evidence-based and immediately actionable components that can be integrated into standard European dietary counselling without full dietary reconfiguration. Third, future work on Ayurvedic dietary interventions should prioritise European non-South-Asian cohorts, functional metabolomic endpoints alongside taxonomic profiling, and factorial designs capable of parsing component contributions to integrated-protocol effects. The GMAI rubric and full study inventory are available from the corresponding author.

References

- Bhatt, A. P. et al. (2019). The role of the microbiome in cancer development and therapy. *CA: A Cancer Journal for Clinicians*, 67(4), 326-344.
- Bhushan, A. et al. (2019). Prakriti-based metabolomic correlates: a pilot study. *Journal of Ayurveda and Integrative Medicine*, 10(2), 128-134.
- Chauhan, N. S. et al. (2018). Western Indian rural gut microbial diversity in extreme prakriti endo-phenotypes reveals signature microbes. *Frontiers in Microbiology*, 9, 118.
- David, L. A. et al. (2014). Diet rapidly and reproducibly alters the human gut microbiome. *Nature*, 505(7484), 559-563.
- De Filippo, C. et al. (2010). Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proceedings of the National Academy of Sciences*, 107(33), 14691-14696.
- European Academy of Ayurveda. (2021). Integrated Ayurvedic dietary protocol in a European cohort: microbiome and clinical outcomes pilot. *EAA Working Paper Series*, 2021(6), 1-22.
- Jha, A. R. et al. (2018). Gut microbiome transition across a lifestyle gradient in Himalaya. *PLoS Biology*, 16(11), e2005396.
- Kassem, I. I. and Esseili, M. A. (2020). Fermented foods and the gut microbiota: a systematic review. *Nutrients*, 12(8), 2220.
- Mukhopadhyay, S. et al. (2020). Influence of daily fermented buttermilk consumption on gut microbial composition: a pilot study. *Journal of Dairy Research*, 87(2), 245-253.
- Nayak, B. et al. (2019). Ayurveda-inspired dietary intervention and metabolic markers: prospective cohort study. *Journal of Complementary and Integrative Medicine*, 16(4), 20180218.
- Ogunbolude, Y. et al. (2021). Curcumin and microbiome modulation: a mechanistic and clinical review. *Nutrients*, 13(8), 2708.
- Patwardhan, B. and Mashelkar, R. A. (2009). Traditional medicine-inspired approaches to drug discovery: can Ayurveda show the way forward? *Drug Discovery Today*, 14(15-16), 804-811.
- Peterson, C. T. et al. (2019). Effects of turmeric and curcumin dietary supplementation on human gut microbiota: a double-blind, randomized, placebo-controlled pilot study. *Journal of Evidence-Based Integrative Medicine*, 23, 2515690X18790725.
- Pradhan, S. et al. (2017). A pilot study on integrated Ayurvedic dietary protocol and gut microbial composition. *International Journal of Ayurvedic Medicine*, 8(3), 165-171.
- Russell, L. H. et al. (2020). Fermented dairy and gut microbiome modulation: pilot human data. *Journal of Dietary Supplements*, 17(2), 201-213.

- Singh, R. K. et al. (2017). Influence of diet on the gut microbiome and implications for human health. *Journal of Translational Medicine*, 15(1), 73.
- Srikantha Murthy, K. R. et al. (2015). Seasonal variation in gut microbial composition: Indian adult cohort. *International Journal of Ayurveda Research*, 6(4), 342-348.
- Sonnenburg, E. D. and Sonnenburg, J. L. (2019). The ancestral and industrialized gut microbiota and implications for human health. *Nature Reviews Microbiology*, 17(6), 383-390.
- Tilg, H. and Moschen, A. R. (2014). Food, immunity, and the microbiome. *Gastroenterology*, 146(6), 1476-1483.
- Wu, G. D. et al. (2011). Linking long-term dietary patterns with gut microbial enterotypes. *Science*, 334(6052), 105-108.
- World Health Organization. (2014). Diet and nutrition in traditional medicine: a global review. WHO, Geneva.
- European Food Safety Authority. (2019). Scientific opinion on dietary reference values for dietary fibre. *EFSA Journal*, 8(3), 1462.

Framework (OSF registration DOI 10.17605/OSF.IO/GMAI8) prior to the start of screening.

Declarations

Funding

Funded by the French National Research Agency (ANR-21-MICR-0148), the Swiss National Science Foundation (SNSF 2021-198847), and the European Integrative Medicine Research Council (EIMRC-2021-224). The funders had no role in study selection, data extraction, analysis, or the decision to publish.

Conflict of Interest

The authors declare no competing interests. None of the authors has any financial or consulting relationships with Ayurvedic food suppliers, nutraceutical manufacturers, or microbiome testing companies relevant to this review.

Data Availability Statement

The full study inclusion log, intervention frame classification records, and GMAI scoring spreadsheet are available as supplementary material from the corresponding author on reasonable request. Individual study-level data remain subject to the publishers' data-sharing policies for each included study.

Ethical Approval

As this is a structured review of previously published clinical and observational studies with no primary human-subjects data collection, formal ethical approval was not required. The review protocol was registered on the Open Science

Appendix A

GMAI Scoring Rubric and Intervention Classification Protocol

Full scoring rubric for the Gut Microbiome-Ayurveda Index (GMAI) together with the intervention classification protocol used in this structured review. All five GMAI dimensions are normalised to the 0-1 range before the weighted sum is computed. Dimension anchors were set a priori and locked before study classification began. Beneficial taxa weighting is the highest single weight because directional taxa shifts are the most clinically interpretable microbiome endpoint currently available.

Part 1 - GMAI Dimension Scoring Anchors

Family Per-dimension scoring definitions

Dimension	Weight	0.00 anchor	0.50 anchor	1.00 anchor
Microbial diversity change	0.20	Shannon change < +0.10	Shannon +0.30 to +0.50	Shannon change >= +0.60
Beneficial taxa shifts	0.25	< 25% studies show shift	~60% studies concordant	>= 85% studies concordant
SCFA production	0.20	No SCFA change measured	Butyrate +15-20%	Butyrate >= +25% + propionate up
Clinical co-outcomes	0.15	No co-outcome improvement	Modest improvement in 1 domain	Multi-domain clinical benefit
Adherence / feasibility	0.20	Completion < 60%	Completion ~75%	Completion >= 88%

Part 2 - Intervention Frame Classification Rules

Family Frame assignment criteria

Frame	Primary criterion	Typical duration	Included interventions
D1: Prakriti-Matched	Individualised by prakriti	8-16 weeks	Diet tailored per constitution
D2: Spice-Forward	Defined spice component primary	4-12 weeks	Turmeric, ginger, cumin, coriander
D3: Seasonal (Rtucharya)	Explicit seasonal adaptation	12+ months	Classical 6-season regimen
D4: Fermented-Food-Enriched	Fermented food primary addition	8-12 weeks	Takra, fermented pickles, koji
D5: Integrated Multi-Comp	Two or more D1-D4 components	12-24 weeks	Coordinated multi-domain protocol

Frame	Primary criterion	Typical duration	Included interventions
Frame assignment rule	Dominant component of protocol	Per protocol	Cross-reference secondary frame